

ON A CONJECTURE OF DYSON

P. M. BLEHER⁽¹⁾ AND P. MAJOR⁽²⁾

⁽¹⁾ Department of Mathematical Sciences,
Indiana University – Purdue University at Indianapolis,
402 N. Blackford Street, Indianapolis, Indiana 46202, USA
E-mail: bleher@math.iupui.edu

⁽²⁾ Mathematical Institute of the Hungarian Academy of Sciences and
Bolyai College of Eötvös Loránd University,
Budapest, HUNGARY
E-mail: major@math-inst.hu

To the memory of Roland L'vovich Dobrushin

ABSTRACT. In this paper we study Dyson's classical r -component hierarchical model with a Hamiltonian function which has a continuous $O(r)$ -symmetry, $r \geq 2$. This is a one-dimensional ferromagnetic model with a long range interaction potential $U(i, j) = -l(d(i, j))d^{-2}(i, j)$, where $d(i, j)$ denotes the hierarchical distance. We are interested in the case when $l(t)$ is a slowly increasing positive function. For a class of free measures, we prove a conjecture of Dyson. This conjecture states that the convergence of the series $l_1 + l_2 + \dots$, where $l_n = l(2^n)$, is a necessary and sufficient condition of the existence of phase transition in the model under consideration, and the spontaneous magnetization vanishes at the critical point, i.e. there is no Thouless' effect. We find however that the distribution of the normalized average spin at the critical temperature T_c tends to the uniform distribution on the unit sphere in $\mathbb{R}^r$ as the volume tends to infinity, a phenomenon which resembles the Thouless effect. We prove that the limit distribution of the average spin is Gaussian for $T > T_c$, and it is non-Gaussian for $T < T_c$. We also show that the density of the limit distribution of the average spin for $T < T_c$ is a nice analytical function which can be found as the unique solution of a nonlinear integral equation. Finally, we determine some critical asymptotics and show that the divergence of the correlation length and magnetic susceptibility is super-polynomial as $T \rightarrow T_c$.

Key words: Dyson's hierarchical model, continuous symmetry, Thouless' effect, renormalization transformation, limit distribution of the average spin, super-polynomial critical asymptotics

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$ - $\text{\texttt{TeX}}$

Contents

1. Introduction. Formulation of the Main Results.
 2. Analytic Reformulation of the Problem. Strategy of the Proof.
 3. Formulation of Auxiliary Theorems.
 4. Basic Estimates in the Low Temperature Region.
 5. Estimates in the Intermediate Region. Proof of Theorem 3.1.
 6. Estimates in the High Temperature Region. Proof of Theorem 3.3.
 7. Estimates in the Low Temperature Region. Proof of Theorem 3.2.
 8. Estimates Near the Critical Point. Proof of Theorems 3.4, 1.3, and 1.5.
- Appendices A and B
- References

1. Introduction. Formulation of the Main Results

In this paper we investigate Dyson's hierarchical vector-valued model with *continuous symmetry*. The model consists of spin variables $\sigma(j) \in \mathbb{R}^r$, $j \in \mathbb{N} = \{1, 2, \dots\}$, where $r \geq 2$. We define the *hierarchical distance* $d(\cdot, \cdot)$ on $\mathbb{N}$ as $d(j, k) = 2^{n(j,k)-1}$ for $j \neq k$, with

$$n(j, k) = \{\min n: \text{there is an integer } l \text{ such that } (l-1)2^n < j, k \leq l2^n\} \quad \text{if } j \neq k,$$

$d(j, j) = 0$. The Hamiltonian of the ferromagnetic hierarchical r -component model in the volume $V_n = \{1, 2, \dots, 2^n\}$ is

$$\mathcal{H}_n(\sigma) = - \sum_{j=1}^{2^n-1} \sum_{k=j+1}^{2^n} \frac{l(d(j, k))}{d^2(j, k)} \sigma(j) \sigma(k), \quad (1.1)$$

where $\sigma(j)\sigma(k)$ denotes a scalar product in $\mathbb{R}^r$, and $l(t)$ is a positive function. In this paper we will be interested in the case when $l(t)$ is a positive increasing function such that

$$\lim_{t \rightarrow \infty} l(t) = \infty; \quad \lim_{t \rightarrow \infty} \frac{l(t)}{t^\varepsilon} = 0, \quad \text{for all } \varepsilon > 0.$$

Let $\nu(dx)$ be a free probability measure on $\mathbb{R}^r$. Then the Gibbs measure in V_n at a temperature $T > 0$ with free boundary conditions is defined as

$$\mu_n(dx; T) = Z_n^{-1}(T) \exp \{-\beta \mathcal{H}_n(x)\} \prod_{j=1}^{2^n} \nu(dx_j), \quad \beta = T^{-1}.$$

We will assume that the free measure $\nu(dx)$ is invariant with respect to the group $O(r)$ of orthogonal transformations, i.e., $\nu(UA) = \nu(A)$ for all $U \in O(r)$ and all Borel sets $A \in B(\mathbb{R}^r)$. Then the Gibbs measure $\mu_n(dx; T)$ is $O(r)$ -invariant as well,

$$\mu_n(UA_1, \dots, UA_{2^n}; T) = \mu_n(A_1, \dots, A_{2^n}; T), \quad \text{for all } U \in O(r), \\ A_j \in B(\mathbb{R}^r), \quad j = 1, \dots, 2^n.$$

In [Dys2], Dyson proved the following theorem (see also [Dys3]). Define

$$l_n = l(2^n). \quad (1.2)$$

Assume that $r = 3$ and $\nu(dx)$ is a uniform measure on the unit sphere in $\mathbb{R}^3$. This is the classical Heisenberg hierarchical model.

Theorem 1.1. (see [Dys2]). *The classical Heisenberg hierarchical model has a phase transition if*

$$B = \sum_{n=1}^{\infty} l_n^{-1} < \infty. \quad (1.3)$$

It has a long-range order so long as

$$\beta > B.$$

Dyson also formulated the following conjecture (see [Dys2]): “It also seems likely that for sequences l_n which are positive and increasing with n the condition (1.3) is necessary for a phase transition in Heisenberg hierarchical models.” The goal of this paper is to *prove* Dyson’s conjecture for a class of hierarchical models and to study the *limit distribution of the average spin* both below and above the critical temperature if condition (1.3) holds. Dyson’s proof is a clever application of *correlation inequalities*. Our approach is based on an analytical study of the *renormalization transformation* for the hierarchical model.

We apply a perturbation technique which only works if the free measure $\nu(dx)$ is a *small perturbation* of the Gaussian measure. Hence, we cannot treat the case when $\nu(dx)$ is a uniform measure on the unit sphere. On the other hand, we will consider *arbitrary* spin dimension $r \geq 2$. We will focus on free measures $\nu(dx)$ which have a density function $p(x)$ on $\mathbb{R}^r$ such that $p(x)$ is close, in an appropriate sense, to the density function

$$p_0(x) = C(\kappa) \exp \left\{ -\frac{|x|^2}{2} - \kappa \frac{|x|^4}{4} \right\} \quad (1.4)$$

with a sufficiently small parameter $\kappa > 0$. Precise conditions on $p(x)$ are given below. We also will assume some regularity conditions about the sequence $l_n = l(2^n)$ (see below).

We are investigating the following question. Let $p_n(x, T)$ denote the density function of the average spin $2^{-n} \sum_{j=1}^{2^n} \sigma(j)$, where $(\sigma(1), \dots, \sigma(2^n))$ is a $\mu_n(T)$ -distributed random vector. Because of the rotational invariance of the model, the function $p_n(x, T)$ is a function of $|x|$. We are interested in the limit behaviour of the function $p_n(x, T)$ as $n \rightarrow \infty$, with an appropriate normalization. In our papers [BM1,3,4] this problem was considered for the polynomial function $l(t) = t^\alpha$ with $0 < \alpha < 1$, when the potential function $l(d(j, k))d^{-2}(j, k)$ in (1.1) is $d^{-2+\alpha}(j, k)$. We have distinguished three cases for α :

- (i) $(1/2) < \alpha < 1$, (ii) $\alpha = 1/2$, and (iii) $0 < \alpha < (1/2)$.

The difference between these cases appears in the asymptotic behavior of $p_n(x, T)$ at small T . When T is small the spontaneous magnetization $M(T)$ is positive, and the function $p_n(x, T)$ is concentrated in a narrow spherical shell near the sphere $|x| = M(T)$. The question is what the width of this shell is and what the limiting shape of $p_n(x, T)$ is like along the radius after an appropriate rescaling. In case (i), the width is of the order of $2^{-n/2}$ and the limit shape of $p_n(x, T)$ is Gaussian (see [BM1]). In case (ii), there is a logarithmic correction in the asymptotics of the width, but the limit shape is still Gaussian (see [BM4]). In case (iii), the width of the shell has a nonstandard asymptotics

of the order of 2^{-n^α} , and the limit shape of $p_n(x, T)$ along the radius (after a rescaling) is a non-Gaussian function which is the solution of a nonlinear integral equation (see [BM3] and the review [BM2]). In the present paper we are interested in the marginal case when $l(t)$ has a sub-polynomial growth.

Before formulating the main results we would like to discuss the importance of Dyson's condition (1.3). In the case of the Ising hierarchical model ($r = 1$), Dyson proved in [Dys2] that there exists the “weakest” interaction function $l(t)$ for which the hierarchical model (1.1) has a phase transition. This function is $l(t) = \log \log t$, which corresponds to $l_n = \log n$. Dyson has proved that if

$$\lim_{n \rightarrow \infty} \frac{l_n}{\log n} = 0, \quad (1.5)$$

then the spontaneous magnetization is equal to zero for all temperatures $T > 0$. On the other hand, if

$$\frac{l_n}{\log n} > \varepsilon \quad \text{for all } n > 0 \text{ with some } \varepsilon > 0,$$

then the spontaneous magnetization is positive at sufficiently low temperatures $T > 0$. In the borderline model, when

$$l_n = J \log n, \quad J > 0, \quad (1.6)$$

Dyson proved that the spontaneous magnetization $M(T)$ has a jump at the critical temperature T_c . The existence of the jump for the 1D Ising model with long-range interaction was first predicted by Thouless (see [Tho], and also the work [AYH] of Anderson, Yuval, and Hamann and references therein) for the translationally invariant Ising model with the interaction

$$H(\sigma) = - \sum_{j,k} \frac{\sigma(j)\sigma(k)}{(j-k)^2}. \quad (1.7)$$

This phenomenon (the jump of $M(T)$ at $T = T_c$) is called the Thouless effect. A rigorous proof of the existence of the Thouless effect in the Ising model with the inverse square interaction (1.7) was given by Aizenman, J. Chayes, L. Chayes, and Newman [ACCN]. Simon proved in [Sim] the absence of continuous symmetry breaking in the one-dimensional r -component model with the interaction (1.7), in the case when $r \geq 2$.

Dyson formulated a general heuristic principle in [Dys2] which tells us when one should expect the Thouless effect in a 1D long-range ferromagnetic model: It should occur for the “weakest” interaction (if it exists) for which a phase transition appears. Dyson wrote that in the hierarchical model “in the Ising case, there exists a borderline model $l_n = \log n$ which is the ‘weakest’ ferromagnet for which a transition occurs, and this borderline model shows a Thouless effect. In the Heisenberg case there exists no borderline model, since there is no ‘most slowly converging’ series (1.3). Thus we

do not expect to find a Thouless effect in any one-dimensional Heisenberg hierarchical ferromagnet.” This conjecture of Dyson, about the absence of a Thouless effect in the Heisenberg case, plays a very essential role in our investigation. We show that in the class of the r -component hierarchical models under consideration, the spontaneous magnetization $M(T)$ approaches zero as T approaches the critical temperature, i.e., there is no Thouless effect. On the other hand, we observe a phenomenon which resembles the Thouless effect: at $T = T_c$ the rescaled distribution

$$\bar{M}_n^r(T_c) p_n(\bar{M}_n(T_c)x, T_c) dx, \quad \bar{M}_n(T) = \left(\int_{\mathbb{R}^r} x^2 p_n(x, T) dx \right)^{1/2}, \quad (1.8)$$

approaches, as $n \rightarrow \infty$, a uniform measure on the unit sphere in $\mathbb{R}^r$, $r \geq 2$. Thus, although the spontaneous magnetization $M(T_c) = \lim_{n \rightarrow \infty} \bar{M}_n(T_c)$ is equal to zero at the critical point, the distribution of the normalized average spin converges to a uniform measure on the unit sphere. This is a “remnant” of the spontaneous magnetization at the critical temperature T_c .

To formulate our results we will need some conditions on the sequence $l_n = l(2^n)$. We need different conditions on l_n in different theorems. We formulate the conditions we shall later apply.

Conditions on the sequence l_n , $n = 0, 1, 2, \dots$: Let us introduce the notation

$$c_n = \frac{l_n}{l_{n-1}}, \quad n = 0, 1, \dots, \text{ with } l_{-1} = 1. \quad (1.9)$$

Condition 1.

$$l_0 = 1; \quad 1 \leq c_n \leq 1.01, \quad \text{for all } n; \quad \lim_{n \rightarrow \infty} c_n = 1. \quad (1.10)$$

Remark. The relation $l_0 = 1$ is not a real condition, it can be reached by a rescaling of the temperature. We use it just for a normalization.

Condition 2.

$$\lim_{n \rightarrow \infty} l_n \sum_{j=n}^{\infty} l_j^{-1} = \infty. \quad (1.11)$$

Moreover, the above relation is uniform in the following sense: For all $\varepsilon > 0$ there are some numbers $K = K(\varepsilon) > 0$ and $L = L(\varepsilon) > 0$ such that

$$l_n \sum_{j=n}^{n+K} l_j^{-1} \geq \varepsilon^{-1} \quad (1.12)$$

for all $n > L$.

Condition 3.

$$\sup_{1 < n < \infty} \sum_{k=1}^n \left(l_k \sum_{j=k}^n l_j^{-1} \right)^{-2} < \infty. \quad (1.13)$$

Condition 4.

$$\sum_{n=1}^{\infty} l_n^{-1} > 400 \kappa^{-1}. \quad (1.14)$$

Condition 5.

$$\frac{l_n}{l_{n+k}} > \bar{\eta} \quad \text{for all } n = 0, 1, 2, \dots, \quad \text{and all } k = 1, \dots, L. \quad (1.15)$$

The numbers $\kappa, \bar{\eta} > 0$, and $L \in \mathbb{N}$ in these conditions will be chosen later. An example of sequences l_n satisfying Conditions 1–5 is given in the following proposition.

Proposition 1.2. *The sequence*

$$l_n = (1 + an)^\lambda, \quad a > 0, \lambda > 1, \quad (1.16)$$

satisfies Conditions 2 and 3 for all $a > 0$ and $\lambda > 1$. There exists a number $a_0 = a_0(\lambda) > 0$ such that this sequence satisfies Condition 1 for all $0 < a < a_0$, a number $a_1 = a_1(\kappa, \lambda) > 0$ such that this sequence satisfies Condition 4 for all $0 < a < a_1$, and finally there exists a number $a_2 = a_2(\bar{\eta}, L) > 0$ such that this sequence satisfies Condition 5 for all $0 < a < a_2$.

Thus, for all $\lambda > 1$ there exists a number

$$a_3 = a_3(\lambda, \kappa, \bar{\eta}, L) = \min\{a_0(\lambda), a_1(\kappa, \lambda), a_2(\bar{\eta}, L)\} > 0$$

such that for all $0 < a < a_3$, the sequence (1.16) satisfy Conditions 1 – 5. We prove Proposition 1.2 in Appendix B below. Now we describe the class of initial densities we shall consider.

Class of initial densities. We say that a probability density $p(x)$ on $\mathbb{R}^r$ belongs to the class $\mathcal{P}_\kappa$ if

$$p(x) = C(1 + \varepsilon(|x|^2)) \exp\left(-\frac{|x|^2}{2} - \kappa \frac{|x|^4}{4}\right), \quad (1.17)$$

where $C > 0$ is a norming factor, and

$$\|\varepsilon(t)\|_{C^4(\mathbb{R}^1)} < 0.01. \quad (1.18)$$

Now we formulate our main results. We denote by $p_n(x, T)$ the distribution of the average spin $2^{-n}[\sigma(1) + \dots + \sigma(2^n)]$ with respect to the Gibbs measure $\mu_n(dx; T)$ and put

$$\bar{M}_n(T) = \left(\int_{\mathbb{R}^r} x^2 p_n(x, T) dx \right)^{1/2} \quad (1.19)$$

By $\bar{p}_n(x, T)$ we denote the rescaled density function

$$\bar{p}_n(x, T) = \bar{M}_n(T) p_n(\bar{M}_n(T)x, T) \quad (1.20)$$

and by $\bar{\nu}_{n,T}(dx)$ the corresponding probability distribution

$$\bar{\nu}_{n,T}(dx) = \bar{p}_n(x, T) dx. \quad (1.21)$$

Formulation of the main results. We fix a sufficiently small positive number η which will be the same through the whole paper. For instance, $\eta = 10^{-100}$ is a good choice. Define the following number $N = N(\eta)$:

$$N = \min\{n: l_n > \eta^{-1}\}. \quad (1.22)$$

Assume that an arbitrary number $\bar{\eta}$ in the interval $0 < \bar{\eta} \leq \eta$ is fixed (it is used in Condition 5).

Theorem 1.3. (Necessity of Dyson's condition). *Assume that*

$$\sum_{n=1}^{\infty} l_n^{-1} = \infty. \quad (1.23)$$

Then there exists a number $\kappa_0 = \kappa_0(N)$ such that for all $0 < \kappa < \kappa_0$ the following statements hold. Assume that the density $p(x) = \frac{\nu(dx)}{dx}$ belongs to the class $\mathcal{P}_\kappa$ and the sequence $\{l_n, n \geq 0\}$ satisfies Conditions 1 — 3. Then there exists a constant $L = L(\bar{\eta}, \kappa)$ such that if the sequence $\{l_n, n \geq 0\}$ satisfies Condition 5, then for all $T > 0$, there exists the limit,

$$\lim_{n \rightarrow \infty} 2^n \bar{M}_n^2(T) = \chi(T) > 0. \quad (1.24)$$

In particular, the spontaneous magnetization satisfies the relation

$$M(T) = \lim_{n \rightarrow \infty} \bar{M}_n(T) = 0. \quad (1.25)$$

In addition, the distribution $\bar{\nu}_{n,T}(dx)$ tends weakly to the standard normal distribution as $n \rightarrow \infty$.

To formulate our results for the case when the Dyson condition (1.3) holds, we define a function $\hat{p}_n(t, T)$ such that

$$p_n(x, T) = \hat{p}_n(|x|, T), \quad (1.26)$$

and introduce the notations

$$V_n(T) = \left(\int_0^\infty (t - \bar{M}_n(T))^2 \hat{p}_n(t, T) dt \right)^{1/2} \quad (1.27)$$

and

$$\pi_n(t, T) = L_n^{-1}(T) \hat{p}_n(\bar{M}_n(T) + V_n(T) t, T), \quad t \geq -\frac{\bar{M}_n(T)}{V_n(T)}, \quad (1.28)$$

where

$$L_n(T) = \int_{-\bar{M}_n(T)/V_n(T)}^\infty \hat{p}_n(\bar{M}_n(T) + V_n(T) t, T) dt.$$

Thus, by (1.26) and (1.28)

$$p_n(x, T) = L_n(T) \pi_n\left(\frac{|x| - \bar{M}_n(T)}{V_n(T)}, T\right). \quad (1.29)$$

Our aim is to prove that in the case when the Dyson condition (1.3) holds, there exists a critical temperature T_c such that the spontaneous magnetization $M(T) = \lim_{n \rightarrow \infty} \bar{M}_n(T)$ is positive for $T < T_c$ and it is zero for $T \geq T_c$. For $T < T_c$ the density function $p_n(x, T)$ is concentrated near a sphere of radius $\bar{M}_n(T)$ and the function $\pi_n(t, T)$ represents a rescaled distribution of $p_n(x, T)$ along the radius $r = |x|$, near the value $r = \bar{M}_n(T)$. We want to prove that $\pi_n(t, T)$ tends to a limit $\pi(t)$ as $n \rightarrow \infty$. It turns out that this limit does exist, and the function $\pi(t)$ is a nice analytic function, although it is non-Gaussian. The function $\pi(t)$ is expressed in terms of a solution of a nonlinear fixed point equation, and the next proposition concerns the existence of such a solution.

Proposition 1.4. *There exists a unique probability density function $g(t)$ on $\mathbb{R}^1$ which satisfies the following fixed point equation:*

$$g(t) = \left(\frac{2}{\sqrt{\pi}} \right)^{r-1} \int_{u \in \mathbb{R}^1, v \in \mathbb{R}^{r-1}} e^{-v^2} g\left(t - u + \frac{v^2}{2}\right) g\left(t + u + \frac{v^2}{2}\right) du dv \quad (1.30)$$

The density $g(t)$ can be extended to an entire function on the complex plane, and for real t it satisfies the estimate

$$0 < g(t) < C_\varepsilon \exp\{-(2 - \varepsilon)|t|\}, \quad \text{for all } \varepsilon > 0. \quad (1.31)$$

For a proof of Proposition 1.4 see the proof of Lemmas 12 and 13 in [BM3].

Theorem 1.5. *Assume that*

$$\sum_{n=1}^\infty l_n^{-1} < \infty.$$

Then there exists a number $\kappa_0 = \kappa_0(N)$ such that for all $0 < \kappa < \kappa_0$ the following statements hold. Assume that the density $p(x) = \frac{\nu(dx)}{dx}$ belongs to the class $\mathcal{P}_\kappa$ and

the sequence $\{l_n, n \geq 0\}$ satisfies Conditions 1 — 4. Then there exists a constant $L = L(\bar{\eta}, \kappa)$ such that if the sequence $\{l_n, n \geq 0\}$ satisfies Condition 5, then there exists a critical temperature $T_c > 0$ with the following properties.

1) If $T > T_c$ then

$$\lim_{n \rightarrow \infty} 2^n \bar{M}_n^2(T) = \chi(T) > 0, \quad (1.32)$$

and the distribution $\bar{\nu}_{n,T}(dx)$ approaches weakly as $n \rightarrow \infty$ a standard normal distribution. The function $\chi(T)$ in (1.32) satisfies the following estimates near the critical point. There exists a temperature $T_0 > T_c$ and numbers $C_2 > C_1 > 0$ such that for all $T_0 > T > T_c$ there exists a number $\bar{n}(T)$ such that

$$\begin{aligned} C_1 \sum_{k=\bar{n}(T)}^{\infty} l_k^{-1} < T - T_c \leq C_2 \sum_{k=\bar{n}(T)}^{\infty} l_k^{-1}, \\ C_1 \frac{2^{\bar{n}(T)}}{l_{\bar{n}(T)}} < \chi(T) < C_2 \frac{2^{\bar{n}(T)}}{l_{\bar{n}(T)}}, \quad \text{for all } T_c < T < T_0. \end{aligned} \quad (1.33)$$

2) If $T = T_c$ then

$$\lim_{n \rightarrow \infty} L_n^{-1} \bar{M}_n(T_c) = 1, \quad (1.34)$$

where

$$L_n = \left(\frac{r-1}{6} \sum_{j=n}^{\infty} l_j^{-1} \right)^{1/2}, \quad (1.35)$$

and the distribution $\bar{\nu}_{n,T_c}(dx)$ tends to the uniform distribution on the unit sphere in $\mathbb{R}^r$ as $n \rightarrow \infty$.

3) If $T < T_c$, then

$$\lim_{n \rightarrow \infty} \bar{M}_n(T) = M(T) > 0, \quad (1.36)$$

and

$$C_1 |T - T_c|^{1/2} < M(T) < C_2 |T - T_c|^{1/2}. \quad (1.37)$$

In addition,

$$\lim_{n \rightarrow \infty} l_n V_n(T) = \gamma(T) \equiv \frac{T}{3M(T)} > 0, \quad (1.38)$$

and

$$\lim_{n \rightarrow \infty} \pi_n(t, T) = \pi(t) \equiv C e^{-2t/3} g(t-a), \quad (1.39)$$

where $g(t)$ is a probability density which satisfies the fixed point equation (1.30), and the quantities C and a are determined from the equations

$$\int_{\mathbb{R}^1} \pi(t) dt = 1, \quad \int_{\mathbb{R}^1} t \pi(t) dt = 0. \quad (1.40)$$

Let us make some remarks about Theorem 1.5. Relations (1.32) and (1.34) imply that

$$M(T) = \lim_{n \rightarrow \infty} \bar{M}_n(T) = 0, \quad \text{for all } T \geq T_c, \quad (1.41)$$

i.e., the spontaneous magnetization $M(T)$ vanishes at $T \geq T_c$. By (1.37),

$$\lim_{T \rightarrow T_c^-} M(T) = 0,$$

hence there is no Thouless effect (by $\lim_{t \rightarrow a^\pm} f(t)$ we denote, as usually, limits of $f(t)$ as $t \rightarrow a$ from the right and from the left, respectively).

The number $\bar{n}(T)$ in (1.33) is very important for our investigation in the subsequent sections. It shows how many iterations of the recursive equation (renormalization group transformation) is needed to reach the “high temperature region” (see Section 3 below for precise definitions). The quantity $\xi(T) = 2^{\bar{n}(T)}$ is the *correlation length*. Usually the correlation length has a power-like asymptotics $\xi(T) \asymp |T - T_c|^{-\nu}$ as $T \rightarrow T_c$ where ν is the critical exponent of the correlation length (see, e.g., [Fi] or [WK]). It follows from (1.33) that in the case under consideration, $\xi(T)$ grows super-polynomially as $T \rightarrow T_c^+$. For instance, if l_n is a sequence determined by equation (1.16) then $\xi(T)$ grows like $\exp [C_0(T - T_c)^{1/(\lambda-1)}]$. Similarly, (1.33) implies that the magnetic susceptibility $\chi(T)$ diverges super-polynomially as $T \rightarrow T_c^+$.

The estimates (1.37) correspond to the value of the critical exponent of spontaneous magnetization $\beta = 1/2$. Relation (1.38) shows that the mean square deviation of the average spin along the radius behaves, when $n \rightarrow \infty$, as

$$V_n(T) \sim \frac{T}{3M(T)l_n}, \quad T < T_c,$$

so that it goes to zero very slowly as $n \rightarrow \infty$ (comparing with the standard behavior of $C2^{-n/2}$). In fact, it goes to zero sub-polynomially with respect to the number of spins 2^n .

Let us say some words about our methods. The questions we investigate in this paper lead to a problem of the following type: We have a starting probability density function $p_0(x, T)$ which depends on a parameter T , the temperature, and we apply the powers of an appropriately defined nonlinear operator $\mathbf{Q}$ to it. This operator $\mathbf{Q}$ is the renormalization group operator. We want to describe the behavior of the sequence of functions $p_n(x, T) = \mathbf{Q}^n p_0(x, T)$, $n = 1, 2, \dots$. In particular, we want to understand how the behavior of this sequence of functions $p_n(x, T)$, $n = 1, 2, \dots$, depends on the parameter T . Our investigation shows that if the function $p_n(x, T)$ is essentially concentrated around the origin, then a negligible error is committed when $p_{n+1}(x, T) = \mathbf{Q}p_n(x, T)$ is replaced by the convolution of the function $p_n(\cdot, T)$ with itself, and this is the case for all n if the parameter T is large. The replacement of the operator $\mathbf{Q}$ by the convolution is called the *high temperature approximation*. On the other hand, if the function $p_n(x, T)$

is essentially concentrated in a narrow shell far from the origin, and this is the case for all n if the parameter T is small, then another good approximation of the function $p_{n+1}(x, T) = \mathbf{O}_n p_n(x, T)$ is possible. This is called the *low temperature approximation*. The high temperature approximation actually means the application of the standard methods of classical probability theory. The low temperature approximation applied in this paper is a natural modification of the methods in our paper [BM3] where a similar problem was investigated. But in the present paper we have to make a more careful and detailed analysis. The reason for it is that while in [BM3] it was enough to investigate only very low temperatures T , now we have to follow carefully when the high and when the low temperature approximation is applicable. Moreover, — and this is a most important part of this paper, — to describe the behavior of the functions $p_n(\cdot, T)$ for all temperatures T we have to follow the behavior of these functions also in the case when neither the high nor the low temperature approximation is applicable. This is the so called *intermediate region*. (See Section 3 for precise definitions).

We study the intermediate region in Section 5. Here we show that if the function $p_n(x, T)$ “is not very far from the origin”, namely, the low temperature approximation is not applicable for it, then the functions $p_{n+k}(x, T)$ are getting closer and closer to the origin as the index $n + k$, $k > 0$ is increasing. Moreover, after finitely many steps k the high temperature approximation is already applicable, and the number of steps we need to get into this situation can be bounded by a constant independent of the parameter T . The proof given in Section 5 contains arguments essentially different from the rest of the paper. Here we heavily exploit that the numbers $c_n = \frac{l_n}{l_{n-1}}$ are very close to one. Informally speaking, the sequence of numbers $c_n - 1$ behaves like a small parameter, and this “small parameter” enables us to handle our model near the critical temperature.

The setup of the rest of the paper is the following. In Section 2 we give an analytic reformulation of the problem and connect Dyson’s condition (1.3) with an approximate recursive formula for some quantities $M_n(T)$ related to the spontaneous magnetization (see (2.28) below). In Section 3 we introduce a notion of low and high temperature regions together with an intermediate region. Then we formulate the basic auxiliary theorems about the characterization of these regions. In Sections 4, 5, and 6 we prove the main estimates concerning the low temperature region, the intermediate region, and the high temperature region, respectively. In Section 7 we prove the convergence of the recursive iterations to the fixed point for all $T < T_c$. Finally, in Section 8 we prove Theorem 3.4 concerning some asymptotics near the critical point T_c and derive Theorems 1.3 and 1.5 from the auxiliary theorems.

2. Analytic Reformulation of the Problem. Strategy of the Proof

The hierarchical structure of the Hamiltonian (1.1) leads to the following *recursive equation* for the density functions $p_n(x, T)$ (see, e.g. Appendix A to the paper [BM2]):

$$p_{n+1}(x, T) = C_n(T) \int \exp\left(\frac{l_n}{T}(x^2 - u^2)\right) p_n(x - u, T) p_n(x + u, T) du, \quad n \geq 0 \quad (2.1)$$

where $p_0(x, T) = p_0(x)$ is defined in (1.17),

$$l_n = l(2^n), \quad (2.2)$$

and $C_n(T)$ is an appropriate norming constant which turns $p_{n+1}(x, T)$ into a density function. We are interested in the asymptotic behaviour of the functions $p_n(x, T)$ as $n \rightarrow \infty$. For the sake of simplicity we will assume that $\varepsilon(t) = 0$ in (1.17), so that $p_0(x)$ coincides with (1.4). All the proofs below are easily extended to the case of nonzero $\varepsilon(t)$ satisfying estimate (1.18).

Put

$$c_n = \frac{l_n}{l_{n-1}}, \quad n = 0, 1, \dots \text{ with } l_{-1} = 1, \quad (2.3)$$

$$A_n = 1 + \sum_{j=1}^{\infty} \frac{c_{n+1}}{2} \dots \frac{c_{n+j}}{2} = 1 + l_n^{-1} \sum_{j=1}^{\infty} 2^{-j} l_{n+j}, \quad n = 0, 1, \dots \quad (2.4)$$

and define

$$\hat{q}_n(x) = \hat{q}_n(x, T) = \exp\left\{\frac{A_n}{2(1+A_n)} l_n x^2\right\} p_n\left(\sqrt{\frac{T}{1+A_n}} x, T\right). \quad (2.5)$$

By (2.3),

$$l_n = \prod_{j=0}^n c_j, \quad n \geq 0, \quad (2.6)$$

by (2.4),

$$l_n A_n = l_n + \frac{l_{n+1} A_{n+1}}{2} \quad (2.7)$$

and from (2.1) we obtain that

$$\hat{q}_{n+1}(x, T) = \bar{C}_n(T) \int e^{-l_n u^2} \hat{q}_n\left(\sqrt{\frac{1+A_n}{1+A_{n+1}}} x - u, T\right) \hat{q}_n\left(\sqrt{\frac{1+A_n}{1+A_{n+1}}} x + u, T\right) du \quad (2.8)$$

with

$$\hat{q}_0(x, T) = C_0(T) \exp\left\{\frac{c_0 A_0 - T}{1+A_0} \frac{x^2}{2} - \frac{\kappa T^2}{(1+A_0)^2} \frac{x^4}{4}\right\}. \quad (2.9)$$

Put

$$\begin{aligned} q_n(x, T) &= (1 + A_n)^{r/2} \hat{q}_n \left(\sqrt{1 + A_n} x, T \right) \\ &= (1 + A_n)^{r/2} \exp \left(\frac{A_n l_n x^2}{2} \right) p_n(\sqrt{T} x, T) \end{aligned} \quad (2.10)$$

and

$$c^{(n)} = (1 + A_{n+1}) l_n, \quad n = 0, 1, 2, \dots \quad (2.11)$$

Then

$$q_{n+1}(x, T) = \frac{1}{Z_n(T)} \int_{\mathbb{R}^r} e^{-c^{(n)} u^2} q_n(x - u, T) q_n(x + u, T) du, \quad (2.12)$$

with

$$q_0(x, T) = \frac{1}{Z_0(T)} \exp \left\{ (c_0 A_0 - T) \frac{x^2}{2} - \kappa T^2 \frac{x^4}{4} \right\}. \quad (2.13)$$

We choose such norming constants in the previous formulas in such a way that

$$\int_{\mathbb{R}^r} q_n(x, T) dx = 1.$$

Thus, the functions $q_n(x, T)$ are defined recursively by formulas (2.12) and (2.13). Our goal is to derive an asymptotics of the functions $q_n(x, T)$ as $n \rightarrow \infty$. Then the asymptotics of the functions $p_n(x, T)$ can be found by means of formula (2.10).

The method of paper [BM3] can be adapted in the study of the low temperature approximation. We shall follow this approach. Due to the rotational symmetry of the Hamiltonian (1.1), the function $q_n(x, T)$ depends only on $|x|$. Define the function $\bar{q}_n(t, T)$, $t \in \mathbb{R}^1$, $n = 0, 1, 2, \dots$, such that

$$q_n(x, T) = C_n^{-1}(T) \bar{q}_n(|x|, T), \quad (2.14)$$

with a norming constant $C_n(T)$ such that $\int_0^\infty \bar{q}_n(t, T) dt = 1$, Put also

$$M_n(T) = \int_0^\infty t \bar{q}_n(t, T) dt, \quad n = 0, 1, \dots, \quad (2.15)$$

and define the functions

$$f_n(t, T) = \frac{1}{c^{(n)}} \bar{q}_n \left(M_n(T) + \frac{t}{c^{(n)}}, T \right), \quad t \in \mathbb{R}^1, \quad n = 0, 1, \dots \quad (2.16)$$

which, as we shall see later, are the appropriate scaling of the functions $\bar{q}_n(t, T)$. Then

$$\bar{q}_n(t, T) = c^{(n)} f_n \left(c^{(n)} (t - M_n(T)), T \right), \quad (2.17)$$

and

$$\int_{-c^{(n)}M_n(T)}^{\infty} f_n(t, T) dt = 1, \quad \int_{-c^{(n)}M_n(T)}^{\infty} t f_n(t, T) dt = 0. \quad (2.18)$$

A low temperature approximation can be applied in the case when $M_n(T)$ is relatively large, comparing with the size of the neighbourhood of $M_n(T)$ in which the function $f_n(t, T)$ is essentially concentrated. In this case we follow the behaviour of the pair $(f_n(t, T), M_n(T))$. To describe this procedure introduce the notation $\mathbf{c} = \{c^{(n)}, n = 0, 1, \dots\}$. The rotational invariance of the function $q_n(\cdot, T)$ suggests the definition of the operator

$$\begin{aligned} \bar{\mathbf{Q}}_{n,M}^{\mathbf{c}} f(t) = \int \exp \left\{ -\frac{u^2}{c^{(n)}} - v^2 \right\} f \left(c^{(n)} \left(\sqrt{\left(M + \frac{t}{c^{(n+1)}} + \frac{u}{c^{(n)}} \right)^2 + \frac{v^2}{c^{(n)}}} - M \right) \right) \\ f \left(c^{(n)} \left(\sqrt{\left(M + \frac{t}{c^{(n+1)}} - \frac{u}{c^{(n)}} \right)^2 + \frac{v^2}{c^{(n)}}} - M \right) \right) du dv. \end{aligned} \quad (2.19)$$

Formula (2.12) together with the definition of the function $f_n(t, T)$ yields that

$$\bar{q}_{n+1} \left(M_n(T) + \frac{t}{c^{(n+1)}}, T \right) = \frac{c^{(n+1)}}{Z_n(T)} \bar{\mathbf{Q}}_{n,M_n(T)}^{\mathbf{c}} f_n(t, T) \quad (2.20)$$

with

$$Z_n(T) = \int_{-c^{(n+1)}M_n(T)}^{\infty} \bar{\mathbf{Q}}_{n,M_n(T)}^{\mathbf{c}} f_n(t, T) dt. \quad (2.21)$$

The norming constant $Z_n(T)$ is determined by the relation $\int_0^{\infty} \bar{q}_{n+1}(t, T) dt = 1$. Define also

$$m_n(T) = m_n(f_n(t, T)) = \frac{1}{Z_n(T)} \int_{-c^{(n+1)}M_n(T)}^{\infty} t \bar{\mathbf{Q}}_{n,M_n(T)}^{\mathbf{c}} f_n(t, T) dt \quad (2.22)$$

and

$$\mathbf{Q}_{n,M_n(T)}^{\mathbf{c}} f_n(t, T) = \frac{1}{Z_n(T)} \bar{\mathbf{Q}}_{n,M_n(T)}^{\mathbf{c}} f_n(t + m_n(T), T). \quad (2.23)$$

Then

$$f_{n+1}(t, T) = \mathbf{Q}_{n,M_n(T)}^{\mathbf{c}} f_n(t, T) \quad \text{and} \quad M_{n+1}(T) = M_n(T) + \frac{m_n(T)}{c^{(n+1)}}. \quad (2.24)$$

The arguments of the function f in the definition of the operator $\bar{\mathbf{Q}}_n$,

$$\ell_{n,M}^{\mathbf{c},\pm}(t, u, v) = c^{(n)} \left(\sqrt{\left(M + \frac{t}{c^{(n+1)}} \pm \frac{u}{c^{(n)}} \right)^2 + \frac{v^2}{c^{(n)}}} - M \right) \quad (2.25)$$

can be well approximated by a simpler expression because of the estimate

$$\left| \ell_{n,M}^{\mathbf{c},\pm}(t,u,v) - \left(\frac{t}{c_{n+1}} \pm u + \frac{v^2}{2M} \right) \right| \leq 100 \left(\frac{v^4}{c^{(n)}M^3} + \frac{t^2 + u^2}{c^{(n)}M} \right) \quad (2.26)$$

which holds for $|t| < \frac{1}{4}c^{(n+1)}M$, $|u| < \frac{1}{4}c^{(n)}M$, and $v^2 < c^{(n)}M^2$. This estimate suggests that for low temperatures T , when $M_n(T)$ is large, the operator $\mathbf{Q}_{n,M_n(T)}^{\mathbf{c}}$ can be well approximated by the operator $\bar{\mathbf{T}}_{n,M_n(T)}^{\mathbf{c}}$ defined as

$$\begin{aligned} \bar{\mathbf{T}}_{n,M_n(T)}^{\mathbf{c}} f(t,T) = \int_{u \in \mathbf{R}^1, v \in \mathbf{R}^{r-1}} e^{-v^2} f \left(\frac{t}{c_{n+1}} + u + \frac{v^2}{2M_n(T)}, T \right) \\ f \left(\frac{t}{c_{n+1}} - u + \frac{v^2}{2M_n(T)}, T \right) du dv \end{aligned} \quad (2.27)$$

The elaboration of the above indicated method will be called the *low temperature approximation*. It works well when $M_n(T)$ is much larger than the range where the function $f_n(t,T)$ is essentially concentrated. For $n=0$ the starting value $M_0(T)$ at low temperatures $T > 0$ is very large. In this case the low temperature expansion can be applied. As we shall see later, the approximation of $\mathbf{Q}_{n,M_n(T)}^{\bar{\mathbf{c}}}$ by $\bar{\mathbf{T}}_{n,M_n(T)}^{\bar{\mathbf{c}}}$ yields that

$$M_{n+1}(T) \sim M_n(T) - \frac{r-1}{4c^{(n)}M_n(T)}, \quad (2.28)$$

which, in turn, implies that

$$M_{n+1}^2(T) \sim M_n^2(T) - \frac{r-1}{2c^{(n)}} \quad (2.29)$$

It follows from (2.4) and (1.10) that

$$2 \leq A_n \leq 2.03, \quad \lim_{n \rightarrow \infty} A_n = 1, \quad (2.30)$$

hence by (2.11),

$$3 \leq \frac{c^{(n)}}{l_n} \leq 3.03, \quad \lim_{n \rightarrow \infty} \frac{c^{(n)}}{l_n} = 3. \quad (2.31)$$

This allows us to rewrite (2.29) as

$$M_{n+1}^2(T) \sim M_n^2(T) - \frac{r-1}{6l_n} \quad (2.32)$$

This formula underlines the importance of the Dyson condition (1.3). Namely, if the series

$$B = \sum_{n=1}^{\infty} l_n^{-1} \quad (2.33)$$

converges then $M_n(T)$ remains large for all n if $T > 0$ is small. Indeed, assume that $T < c_0 A_0/2$. Then it follows from (2.13) that $M_0^2(T) > C(\kappa T^2)^{-1}$, hence by (2.32),

$$M_n^2(T) \geq M_0^2(T) - \frac{r-1}{6} \sum_{n=0}^{\infty} l_n^{-1} \geq C(\kappa T^2)^{-1} - C_1 \gg 1$$

for all n if $T > 0$ is small, which was stated. On the other hand, if the series (2.33) diverges, then for some n , $M_n(T)$ becomes small, and the approximation (2.28) becomes inapplicable.

The low temperature approximation can be applied when $M_n(T)$ is large. When $M_n(T)$ is small a different approximation is natural. If the function $q_n(x, T)$ is essentially concentrated in a ball whose radius is much less than $(c^{(n)})^{-1/2}$, then a small error is committed if the kernel function $e^{-c^{(n)}u^2}$ in formula (2.12) is omitted. This means that the formula expressing $q_{n+1}(x)$ by $q_n(x)$ can be well approximated through the convolution $q_{n+1}(2x) = q_n * q_n(2x)$. This approximation will be called the *high temperature approximation*. If the high temperature approximation can be applied for $q_n(x, T)$, then the function $q_{n+1}(x, T)$ is even more strongly concentrated around zero. Hence, as a detailed analysis will show, if at a temperature T it can be applied for a certain n_0 , then it can be applied for all $n \geq n_0$.

Finally, there are such pairs (n, T) for which the function $f_n(x, T)$ can be studied neither by the low nor by the high temperature approximation. We call the set of such pairs an *intermediate region*. We shall prove that if the sequence $c^{(n)}$ sufficiently slowly tends to infinity and the function $f_n(x, T)$ is out of the region where the low temperature approximation is applicable, then the density function $f_{n+1}(x, T)$ will be more strongly concentrated around zero than the function $f_n(x, T)$. Moreover, in *finitely many* steps the function $f_{n+k}(x, T)$ will be so strongly concentrated around zero that after this step the high temperature approximation is applicable. It is important that the number of steps k needed to get into the high temperature region can be bounded independently of the parameter T .

The main part of the paper consists of an elaboration of the above heuristic argument.

3. Formulation of Auxiliary Theorems

To describe the region where the low temperature approximation will be applied we define some sequences $\beta_n(T)$ which depend on the temperature T . Define recursively,

$$\begin{aligned}\beta_N(T) &= \frac{(c^{(N)})^2}{2^N}, \\ \beta_{n+1}(T) &= \left(\frac{c_{n+1}^2}{2} + \sqrt{\frac{\beta_n(T)}{c^{(n)}}} \right) \beta_n(T) + \frac{10}{M_n^2(T)} \quad \text{for } n > N,\end{aligned}\tag{3.1}$$

where the number N is defined in (1.22) and $M_n(T)$ in (2.15). As it will be seen later, these numbers measure how strongly the functions $f_n(x, T)$ are concentrated around zero. We define the low temperature region, where low temperature approximation will be applied.

Definition of the low temperature region. *A pair (n, T) is in the low temperature region if $0 < T \leq c_0 A_0/2$, where A_0 defined in (2.4), and either $0 \leq n \leq N$ with the number N defined in (1.22) or $n > N$ and $\frac{\beta_{n-1}(T)}{c^{(n-1)}} \leq \eta$. The temperature T is in the low temperature region if the pair (n, T) is in the low temperature region for all numbers n .*

Let us remark that by (2.6) and (1.10)

$$1 \leq l_n = \prod_{j=1}^n c_j \leq 1.01^n,\tag{3.2}$$

hence by (2.31),

$$3 \leq c^{(n)} \leq 3.03 \cdot 1.01^n.\tag{3.3}$$

Therefore, by (3.1),

$$\frac{\beta_N(T)}{c^{(N)}} = \frac{c^{(N)}}{2^N} \leq \frac{1}{c^{(N)}} \leq \eta\tag{3.4}$$

hence the pair $(N+1, T)$ is in the low temperature region if $T \leq c_0 A_0/2$. Since $\beta_{n+1}(T) \geq \frac{10}{M_n^2(T)}$ the pair (n, T) can get out of the low temperature region only if $M_n(T)$ becomes very small.

To define the high temperature region introduce the notations

$$\begin{aligned}h_n(x, T) &= \left(c^{(n)}\right)^{-r/2} q_n\left(\frac{x}{\sqrt{c^{(n)}}}, T\right), \\ D_n^2(T) &= \int_{\mathbb{R}^r} x^2 h_n(x, T) dx.\end{aligned}\tag{3.5}$$

where the function $q_n(x, T)$ is defined in (2.10). Let us also introduce the probability measure $H_{n,T}$,

$$H_{n,T}(\mathbf{A}) = \int_{\mathbf{A}} h_n(x, T) dx, \quad \mathbf{A} \subset \mathbb{R}^r\tag{3.6}$$

on $\mathbb{R}^r$.

Definition of the high temperature region. A pair (n, T) is in the high temperature region if $D_n^2(T) < e^{-1/\eta^2}$, where $D_n^2(T)$ is defined in (3.5). The temperature T is in the high temperature region if there exists a threshold index $n_0(T)$ such that (n, T) is in the high temperature region for all $n \geq n_0(T)$.

It may happen that a pair (n, T) belongs neither to the low nor to the high temperature region. Then we say that (n, T) belongs to the *intermediate region*. The following result is very important for us.

Theorem 3.1. *There exists a number $\kappa_0 = \kappa_0(N)$ such that for all $0 < \kappa < \kappa_0$ there exists $L = L(\bar{\eta}, \kappa)$ such that the following is true. Assume that Conditions 1 and 5 hold, and that for a temperature $T > 0$, there exist pairs (n, T) which do not belong to the low temperature region. Let $\bar{n}(T) \geq 0$ be the smallest such number.*

Assume that the pair $(\bar{n}(T), T)$ does not belong to the high temperature region. (In this case $(\bar{n}(T), T)$ is in the intermediate region.) Then there exist numbers $K = K(\bar{\eta}, t) > 0$, $\tilde{\eta} = \tilde{\eta}(\bar{\eta}, t)$, and $k = k(\bar{\eta}, t) \in \mathbb{N}$ such that

$$D_{\bar{n}(T)}^2(T) < K, \quad \tilde{\eta} < D_{\bar{n}(T)+k}^2(T) < e^{-1/\eta^2}, \quad (3.7)$$

which implies in particular that the pair $(\bar{n}(T) + k, T)$ with this index k belongs to the high temperature region.

Corollary. *Under the conditions of Theorem 3.1 all temperatures $T > 0$ belong to either the low or the high temperature region. If the Dyson condition (1.3) holds, then all sufficiently low temperatures belong to the low and all sufficiently high temperatures to the high temperature region. If the Dyson condition (1.3) is violated, then all temperatures $T > 0$ belong to the high temperature region.*

Next theorem concerns the low temperature region.

Theorem 3.2. *There exists a number $\kappa_0 = \kappa_0(N)$ such that for all $0 < \kappa < \kappa_0$ the following is true. Assume that the Dyson condition (1.3) and Conditions 1 and 2 hold. Assume that the temperature T is in the low temperature region. Then the limit*

$$\lim_{n \rightarrow \infty} M_n(T) = M_\infty(T) \quad (3.8)$$

exists, and

$$\lim_{n \rightarrow \infty} \frac{M_n^2(T) - M_\infty^2(T)}{\frac{r-1}{2} \sum_{k=n}^{\infty} \frac{1}{c^{(k)}}} = 1. \quad (3.9)$$

In addition,

$$\lim_{n \rightarrow \infty} \left\| \frac{1}{M_n(T)} f_n \left(\frac{t}{M_n(T)}, T \right) - g \left(t - \frac{r-1}{4} \right) \right\| = 0, \quad (3.10)$$

where

$$\|f(t)\| = \sum_{j=0}^2 \sup_{t \geq -c^{(n)} M_n^2(T)} e^{|t|} \left| \frac{d^j f(t)}{dt^j} \right| \quad (3.11)$$

and the probability density $g(t)$ is defined as a solution of the fixed point equation (1.30).

Remark. Observe that the value $\frac{r-1}{4}$ of the shift of the function $g(t)$ in (3.10) fits the equation

$$\int_{\mathbb{R}^1} t g\left(t - \frac{r-1}{4}\right) dt = 0.$$

From this theorem, the Part 3) of Theorem 1.5 follows, with the exception of estimate (1.37). Indeed, we can express the function $p_n(x, T)$ in terms of $f_n(t, T)$. Namely, by (2.10), (2.14), and (2.17),

$$p_n(x, T) = L_n^{-1}(T) \exp\left(-\frac{A_n l_n |x|^2}{2T}\right) f_n\left(\frac{c^{(n)}}{\sqrt{T}}\left(|x| - \sqrt{T} M_n(T)\right), T\right) \quad (3.12)$$

Let us write that $|x|^2 = \left(\sqrt{T} M_n(T) + |x| - \sqrt{T} M_n(T)\right)^2$, hence

$$\begin{aligned} \exp\left(-\frac{A_n l_n |x|^2}{2T}\right) &= \exp\left\{-\frac{A_n l_n}{2T} [T M_n^2(T) + 2\sqrt{T} M_n(T)(|x| - \sqrt{T} M_n(T)) \right. \\ &\quad \left. + (|x| - \sqrt{T} M_n(T))^2]\right\}, \end{aligned} \quad (3.13)$$

and substitute it into (3.12). This leads to the equation

$$p_n(x, T) = \tilde{L}_n^{-1}(T) \tilde{f}_n\left(\frac{|x| - \tilde{M}_n(T)}{\tilde{V}_n(T)}, T\right), \quad (3.14)$$

where

$$\begin{aligned} \tilde{M}_n(T) &= \sqrt{T} M_n(T), \quad \tilde{V}_n(T) = \frac{\sqrt{T}}{c^{(n)} M_n(T)} \\ \tilde{f}_n(t, T) &= f_n\left(\frac{t}{M_n(T)}, T\right) \exp\left(-\frac{A_n l_n t}{c^{(n)}} - \varepsilon_n(t, T)\right) \\ \varepsilon_n(t, T) &= \frac{A_n l_n t^2}{2(c^{(n)})^2 M_n^2(T)}. \end{aligned} \quad (3.15)$$

Observe that by (2.30) and (2.31)

$$\lim_{n \rightarrow \infty} \frac{A_n l_n}{c^{(n)}} = \frac{2}{3}, \quad \lim_{n \rightarrow \infty} \frac{A_n l_n}{2(c^{(n)})^2 M_n^2(T)} = 0, \quad (3.16)$$

hence (3.10) implies that there is some $C_0 > 0$ such that

$$\lim_{n \rightarrow \infty} \left\| \tilde{f}_n(t, T) - C_0 g\left(t - \frac{r-1}{4}\right) e^{-2t/3} \right\|' = 0, \quad (3.17)$$

where

$$\|f(t)\|' = \sum_{j=0}^2 \sup_{t \geq -c^{(n)} M_n^2(T)} e^{|t|/3} \left| \frac{d^j f(t)}{dt^j} \right|. \quad (3.18)$$

It remains to shift $\tilde{f}_n(t)$ to secure the mean value to be zero. Consider $\pi(t)$ defined in (1.31). Put

$$\tilde{\pi}_n(t, T) = C_n(T) \tilde{f}_n(t - a_0, T), \quad a_0 = a - \frac{r-1}{4}, \quad (3.19)$$

where $C_n(T)$ is a norming factor such that

$$\int_{\mathbb{R}^1} \tilde{\pi}_n(t) dt = 1. \quad (3.20)$$

Then

$$p_n(x, T) = L_n^{-1}(T) \tilde{\pi}_n \left(\frac{|x| - M_{n0}(T)}{\tilde{V}_n(T)}, T \right) \quad \text{with } M_{n0}(T) = \tilde{M}_n + a_0 \tilde{V}_n(T) \quad (3.21)$$

and

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}^1} t \pi_n(t) dt = 0. \quad (3.22)$$

Comparing this formula with (1.29) we obtain (1.36), (1.38), and (1.39) with

$$M(T) = \sqrt{T} M_\infty(T), \quad \gamma(T) = \frac{\sqrt{T}}{3M_\infty(T)} = \frac{T}{3M(T)}, \quad (3.23)$$

where $M_\infty(T)$ is the limit (3.4).

Now we formulate a theorem about the *high temperature region*. Put

$$\tilde{h}_n(x, T) = 2^{-rn/2} q_n \left(2^{-n/2} x, T \right) = \left(\frac{c^{(n)}}{2^n} \right)^{r/2} h_n \left(\sqrt{\frac{c^{(n)}}{2^n}} x, T \right), \quad (3.24)$$

and define the probability measures

$$\tilde{H}_{n,T}(\mathbf{A}) = \int_{\mathbf{A}} \tilde{h}_n(x, T) dx, \quad \mathbf{A} \subset \mathbb{R}^r \quad (3.25)$$

on $\mathbb{R}^r$.

Theorem 3.3. *There exists a number $\kappa_0 = \kappa_0(N)$ such that for all $0 < \kappa < \kappa_0$ there exists a number $L = L(\bar{\eta}, \kappa)$ such that the following is true. Assume that Conditions 1 and 5 hold and that T is in the high temperature region. Then the measures $\tilde{H}_{n,T}$ defined in (3.25) converge weakly to a normal distribution on $\mathbb{R}^r$ with expectation zero and variance $\sigma^2(T)\mathbf{I}$ with some $\sigma^2(T) > 0$, where $\mathbf{I}$ denotes the identity matrix.*

If T belongs to the high temperature region, but the pair $n = (0, T)$ does not belong to it, (i.e. the temperature T is not too high) then the inequality

$$C_1 \frac{2^{\bar{n}(T)}}{c^{\bar{n}(T)}} \leq \sigma^2(T) \leq C_2 \frac{2^{\bar{n}(T)}}{c^{\bar{n}(T)}} \quad (3.26)$$

also holds with some $C_2 > C_1 > 0$, where $\bar{n}(T)$ is defined in Theorem 3.1.

Remark. Not only the convergence of the measures $\tilde{H}_{n,T}$ but also the convergence of their density functions $\tilde{h}_n(x, T)$ could be proved. But the proof of the convergence of the distribution is simpler, and it is also sufficient for our purposes.

Corollary. Let $\bar{H}_{n,T}$ denote the probability measure on $\mathbb{R}^r$ with the density function $2^{-rn/2} T^r p_n(2^{-n/2} \sqrt{T}x, T)$. Under the conditions of Theorem 3.3 the measures $\bar{H}_{n,T}$ have the same Gaussian limit as the measures $\tilde{H}_{n,T}$ defined in Theorem 3.3 as $n \rightarrow \infty$.

Our last theorem concerns the *critical point*. We want to show that there is a critical temperature $T_{\text{cr.}}$ above which all temperatures belong to the high and below which all temperatures belong to the low temperature region. We also want to describe the situation in the neighbourhood of the critical temperature in more detail. In Theorem 3.4 we prove such a result.

Theorem 3.4. There exists a number $\kappa_0 = \kappa_0(N)$ such that for all $0 < \kappa < \kappa_0$ there exists a number $L = L(\bar{\eta}, \kappa)$ such that the following is true. Assume that Conditions 1 — 4 are satisfied. Then for a fixed n the set of temperatures T for which (n, T) belongs to the low temperature region forms an interval $(0, T_n]$, and the sequence T_n , $n = 1, 2, \dots$, is monotone decreasing in n . Define the critical temperature $T_{\text{cr.}}$ as the limit $T_{\text{cr.}} = \lim_{n \rightarrow \infty} T_n$. Then $c_0 A_0 / 4 > T_{\text{cr.}} > 0$. The function $M_\infty(T) = \lim_{n \rightarrow \infty} M_n(T)$ exists in the interval $(0, T_{\text{cr.}}]$, and for fixed n the function $M_n(\cdot)$ is strictly decreasing in the interval $(0, T_n]$. The relation $M_\infty(T_{\text{cr.}}) = 0$ holds. If $T_{\text{cr.}} + \varepsilon > T > T_{\text{cr.}}$ with some $\varepsilon > 0$, then the inequality

$$C_1 \sum_{k=\bar{n}(T)}^{\infty} \frac{1}{c^{(k)}} < T - T_{\text{cr.}} < C_2 \sum_{k=\bar{n}(T)}^{\infty} \frac{1}{c^{(k)}} \quad (3.27)$$

holds with some appropriate numbers $C_2 > C_1 > 0$, where $\bar{n}(T)$ is defined in Theorem 3.1. If $T_{\text{cr.}} - \varepsilon < T < T_{\text{cr.}}$ with a sufficiently small $\varepsilon > 0$, then

$$C_1 (T_{\text{cr.}} - T)^{1/2} < M_\infty(T) < C_2 (T_{\text{cr.}} - T)^{1/2}. \quad (3.28)$$

4. Basic Estimates in the Low Temperature Region

In this section we give some basic estimates on the function $f_n(x, T)$ and its derivatives (with respect to the variable x) if the pair (n, T) is in the low temperature region. These estimates state, in particular, that in the definition of the functions $f_n(x, T)$ the right scaling was chosen. With the scaling in formula (2.16) the function $f_n(x, T)$ is essentially concentrated in a finite interval whose size depends only on $M_n(T)$.

Both the results and proofs are closely related to those of Sections 3 — 6 in paper [BM3]. For the sake of simpler notations we shall assume that $\mathbb{R}^r = \mathbb{R}^2$, i.e., we work in two dimensional models. But all proofs can be simply generalized to the case $r \geq 2$. To simplify notations further, in this section we will denote the restriction of the function $q_n(x, T)$ defined in (2.10) (with $x \in \mathbb{R}^r$) to the ray $l = \{x = (x_1, 0, \dots, 0), x_1 \geq 0\}$. again by $q_n(x, T), x \in \mathbb{R}^1, x \geq 0$. Since the function $q_n(x, T)$ in (2.10) depends only on $|x|$, this restriction determines the original function $q_n(\cdot, T)$ uniquely.

First we consider the case of *small* indices $0 \leq n \leq N$, where the number N defined in (1.22) (cf. Section 4 in [BM3]), and we begin with $n = 0$. Assume that $T < c_0 A_0/2$ and $\kappa > 0$ is small (exact conditions on the smallness of κ will be given later). In this case the function $q_0(x, T)$ has its maximum in the points $\pm \hat{M}_0(T)$ (see (2.13)) where

$$\hat{M}_0(T) = \left(\frac{A_0 c_0 - T}{\kappa T^2} \right). \quad (4.1)$$

is a large number. From (2.13) we obtain that

$$\frac{1}{c^{(0)}} q_0 \left(\hat{M}_0(T) + \frac{x}{c^{(0)}}, T \right) = \frac{1}{Z_0(T)} \exp \left\{ - (A_0 c_0 - T) \left(\frac{x}{c^{(0)}} \right)^2 \left(1 + \frac{x}{2c^{(0)} \hat{M}_0(T)} \right)^2 \right\} \quad (4.2)$$

where

$$Z_0(T) = \int_{-\hat{M}_0(T)}^{\infty} \exp \left\{ - (A_0 c_0 - T) \left(\frac{x}{c^{(0)}} \right)^2 \left(1 + \frac{x}{2c^{(0)} \hat{M}_0(T)} \right)^2 \right\} dx. \quad (4.2')$$

It can be proved by means of the identity

$$M_0(T) - \hat{M}_0(T) = \frac{\int_{-\hat{M}_0(T)}^{\infty} x \exp \left\{ - (c_0 A_0 - T) \left(\frac{x}{c^{(0)}} \right)^2 \left(1 + \frac{x}{2c^{(0)} \hat{M}_0(T)} \right)^2 \right\} dx}{\int_{-\hat{M}_0(T)}^{\infty} \exp \left\{ - (A_0 c_0 - T) \left(\frac{x}{c^{(0)}} \right)^2 \left(1 + \frac{x}{2c^{(0)} \hat{M}_0(T)} \right)^2 \right\} dx} \quad (4.3)$$

that

$$\left| M_0(T) - \hat{M}_0(T) \right| \leq \frac{\text{const.}}{M_0(T)} \leq \text{const.} \sqrt{\kappa} T \quad (4.4)$$

where $M_0(T)$ is defined (2.15). This shows that $\hat{M}_0(T)$ is a very good approximation to $M_0(T)$. Some calculation yields, with the help of formulas (4.1) and (4.3), that

$$\left| Z_0(T) - \frac{c^{(0)}\sqrt{\pi}}{\sqrt{(A_0c_0 - T)}} \right| \leq \text{const.} \sqrt{\kappa} T \quad (4.5)$$

and from (4.1)–(4.5) we obtain that

$$\left| \frac{\partial^j}{\partial x^j} \left(f_0(x, T) - \frac{\sqrt{A_0c_0 - T}}{c^{(0)}\sqrt{\pi}} \exp \left\{ -(A_0c_0 - T) \left(\frac{x}{c^{(0)}} \right)^2 \right\} \right) \right| \leq \text{const.} \kappa^{1/4} e^{-2|x|/c^{(0)}} \quad (4.6)$$

if $|x| < \log \kappa^{-1}$, $j = 0, 1, 2$,

and

$$\left| \frac{\partial^j}{\partial x^j} f_0(x, T) \right| \leq C \exp \left\{ -\frac{(A_0c_0 - T)}{4c^{(0)}} \left| 2x + \frac{x^2}{c^{(0)}M_0^2(T)} \right| \right\} \quad (4.7)$$

for $x \geq -c^{(0)}M_0(T)$, $j = 0, 1, 2$.

A relatively small error is committed if M_n is very large and the arguments $\ell_{n, M_n}^\pm(x, u, v)$ (defined in formula (2.9)) of the function f_n in the operator $\bar{\mathbf{Q}}_{n, M}^{\mathbf{c}} f_n$ are replaced by $x \pm u$. Exploiting this fact one can prove, using a natural adaptation of the proof of Proposition 1 of paper [BM3], the following

Proposition 4.1. *There exists a number $\kappa_0 = \kappa_0(N)$ such that if (i) $0 < \kappa < \kappa_0$, (ii) $0 < T \leq c_0 A_0/2$, and (iii) Condition 1 holds, then the relations*

$$\left| \frac{\partial^j}{\partial x^j} \left(f_n(x, T) - \frac{\sqrt{A_0c_0 - T}}{\sqrt{\pi}} \frac{2^n}{c^{(n)}} \exp \left\{ -2^n (A_0c_0 - T) \left(\frac{x}{c^{(n)}} \right)^2 \right\} \right) \right| \leq B(n) \kappa^{1/4} e^{-2^{n+1}|x|/c^{(n)}}, \quad \text{if } |x| < 2^{-n} \log \kappa^{-1}, \quad j = 0, 1, 2, \quad (4.8)$$

$$\left| \frac{\partial^j}{\partial x^j} f_n(x, T) \right| \leq B(n) \exp \left\{ -\frac{(A_0c_0 - T)}{4} \frac{2^n}{c^{(n)}} \left| 2x + \frac{x^2}{c^{(n)}M_n^2(T)} \right| \right\} \quad (4.9)$$

for $x \geq -c^{(n)}M_n(T)$, $j = 0, 1, 2$,

and

$$|M_n(T) - \hat{M}_0(T)| \leq B(n) \sqrt{\kappa} T \quad (4.10)$$

hold for all $0 \leq n \leq N$ with the function $\hat{M}_0(T)$ defined in (4.1) and a function $B(n)$ which depends neither on T nor κ .

We formulate and prove, similarly to paper [BM3], certain inductive hypotheses about the behaviour of the functions $f_n(x, T)$ for $n \geq N$ if the pair (n, T) in the low

temperature region. In the formulation of these hypotheses we apply the sequence $\beta_n(T)$ defined in (3.1) and the sequence $\alpha_n(T)$ defined as

$$\begin{aligned}\alpha_N(T) &= \frac{1}{200} \frac{c^{(N)^2}}{2^N} \\ \alpha_{n+1}(T) &= \left(\frac{c_{n+1}^2}{2} - \sqrt{\frac{\beta_n(T)}{c^{(n)}}} \right) \alpha_n(T) + \frac{10^{-12}}{M_n^2(T)} \quad \text{for } n > N\end{aligned}\tag{4.11}$$

To formulate the inductive hypotheses we also introduce a regularization of the functions $f_n(x, T)$.

Definition of the regularization of the functions $f_n(x, T)$. *Let us fix a C^∞ -function $\varphi(x)$, $-\infty < x < \infty$, such that $\varphi(x) = 1$ for $|x| \leq 1$, $0 \leq \varphi(x) \leq 1$ if $1 \leq x \leq 2$ and $\varphi(x) = 0$ for $|x| \geq 2$. Then the regularization of the function $f_n(x, T)$ is $\varphi_n(f_n(x, T)) = A_n \varphi\left(\frac{x + B_n}{100\sqrt{c^{(n)}}}\right) f_n(x + B_n, T)$ with such norming constants A_n and B_n for which $\int_{-\infty}^{\infty} \varphi_n(f_n(x, T)) dx = 1$, and $\int_{-\infty}^{\infty} x \varphi_n(f_n(x, T)) dx = 0$.*

Now we formulate the inductive hypotheses.

Hypothesis $I(n)$.

$$\begin{aligned}\left| \frac{\partial^j f_n(x, T)}{\partial x^j} \right| &\leq \frac{C}{\beta_n(T)^{(j+1)/2}} \exp \left\{ -\frac{1}{\sqrt{\beta_n(T)}} \left| 2x + \frac{x^2}{c^{(n)} M_n(T)} \right| \right\} \\ &\quad \text{for } j = 0, 1, 2, \quad x \geq -c^{(n)} M_n(T).\end{aligned}$$

with a universal constant $C > 0$. One could choose e.g. $C = 10^{20}$.

Hypothesis $J(n)$.

$$|\tilde{\varphi}_n f_n(t + is, T)| \leq \frac{e^{\beta_n(T)s^2}}{1 + \alpha_n(T)t^2} \quad \text{if } |s| \leq \frac{2}{\sqrt{\beta_{n+1}(T)}}$$

with $\tilde{\varphi}_n f_n(t + is, T) = \int e^{(it-s)x} f_n(x, T) dx$, i.e. it is the Fourier transform (with a different norming constant) of the function $f_n(x, T)$ together with its analytic continuation.

We need Proposition 4.1 because of its consequence formulated below. Its proof can also be found in [BM3].

Corollary of Proposition 4.1. *Under the conditions of Proposition 4.1 the inductive hypotheses $I(n)$ and $J(n)$ hold for $n = N$ with a universal constant $C > 0$ in hypothesis $I(n)$. (For instance one can choose $C = 10^5$.)*

Before formulating the main result of this Section we introduce the operators $\mathbf{T}_n$. They are appropriate scaling of the operators $\bar{\mathbf{T}}_{n, M_n(T)}^c$ defined in formula (3.11), but

these operators will be applied only for the regularization of the functions $f_n(x, T)$ and not for the functions $f_n(x, T)$ themselves. Put

$$\begin{aligned} \mathbf{T}_n \varphi_n(f_n(x, T)) &= \frac{2}{c_{n+1} \sqrt{\pi}} \int e^{-v^2} \varphi_n \left(f_n \left(\frac{x}{c_{n+1}} - \frac{1}{4M_n(T)} + u + \frac{v^2}{2M_n(T)}, T \right) \right) \\ &\quad \varphi_n \left(f_n \left(\frac{x}{c_{n+1}} - \frac{1}{4M_n(T)} - u + \frac{v^2}{2M_n(T)}, T \right) \right) du dv. \end{aligned} \quad (4.12)$$

The main result of this section is the following

Proposition 4.2. *There exists $\kappa_0 = \kappa_0(N) > 0$ such that if (i) $0 < \kappa < \kappa_0$, (in formula (1.4)) (ii) Condition 1 holds, and (iii) the pairs (m, T) belong to the low domain region for all $0 \leq m \leq n$, then the function $f_{n+1}(x, T)$ satisfies the inductive hypotheses $I(n+1)$ and $J(n+1)$ with the same universal constant $C > 0$ (independent of κ , n , η and T). Also the relation*

$$M_{n+1}(T) = M_n(T) - \frac{1}{4c^{(n)}M_n(T)} + \frac{\gamma_n(T)}{c^{(n)}} \quad \text{with } |\gamma_n(T)| \leq C_1 \frac{\beta_{n+1}(T)}{c^{(n+1)}} \sqrt{\beta_{n+1}(T)} \quad (4.13)$$

holds with a universal constant $C_1 > 0$ together with the inequalities

$$1 \leq \frac{\beta_{n+1}(T)}{\alpha_{n+1}(T)} \leq K_1, \quad (4.14)$$

$$\begin{aligned} \left| \frac{\partial^j}{\partial x^j} (f_{n+1}(x, T) - \mathbf{T}_n \varphi_n(f_n(x, T))) \right| &\leq \frac{K_2 C^4}{\beta_{n+1}^{(j+1)/2}(T)} \frac{\beta_n(T)}{c^{(n)}} \\ &\quad \left[\exp \left\{ -\frac{1}{\sqrt{\beta_{n+1}(T)}} \left| 2x + \frac{x^2}{c^{(n+1)}M_{n+1}(T)} \right| \right\} + \exp \left\{ -\frac{2|x|}{\sqrt{\beta_{n+1}(T)}} \right\} \right] \\ &\quad x > -c^{(n+1)}M_{n+1}(T), \quad j = 0, 1, 2 \end{aligned} \quad (4.15)$$

and

$$\begin{aligned} \left| \frac{\partial^j}{\partial x^j} \mathbf{T}_n \varphi_n(f_n(x, T)) \right| &\leq \frac{K_3 C^2}{\beta_{n+1}^{(j+1)/2}(T)} \exp \left\{ -\frac{2|x|}{\sqrt{\beta_{n+1}(T)}} \right\}, \\ &\quad x \in \mathbb{R}^1, \quad j = 0, 1, 2, 3, 4 \end{aligned} \quad (4.16)$$

with some universal constants K_1 , K_2 and K_3 .

The proof of Proposition 4.2 is based on the observation that the operator $\mathbf{T}_n$ approximates the operator $\mathbf{Q}_{n, M_n(T)}^c$ very well, it has a relatively simple structure, it is actually a convolution. More explicitly, it can be written in the form

$$\begin{aligned} \mathbf{T}_n \varphi_n(f_n(x, T)) &= \frac{4}{c_{n+1} \sqrt{\pi}} \int_0^\infty e^{-v^2} \varphi_n(f_n) * \varphi_n(f_n) \left(\frac{2x}{c_{n+1}} + \frac{v^2}{M_n(T)} - \frac{1}{2M_n(T)}, T \right) \\ &= \frac{2}{c_{n+1}} \varphi_n(f_n) * \varphi_n(f_n) * k_{M_n(T)} \left(\frac{2x}{c_{n+1}} \right), \end{aligned}$$

where $*$ denotes convolution, and $k_{M_n(T)}(x) = M_n(T)k(M_n(T)x)$ with $k(x) = \frac{1}{\sqrt{\pi x}}e^{-x}$ for $x > 0$ and $k(x) = 0$ for $x \leq 0$.

The operator $\mathbf{T}_n$ has a certain contraction property which can be expressed in the Fourier space. The Fourier transform of $\tilde{\mathbf{T}}_n \tilde{\varphi}_n(f_n(\xi, T))$ can be expressed as

$$\tilde{\mathbf{T}}_n \tilde{\varphi}_n(f_n(\xi, T)) = \frac{\exp \left\{ i \frac{c_{n+1}}{4M_n(T)} \xi \right\}}{\sqrt{1 + i \frac{c_{n+1}}{2M_n(T)} \xi}} \tilde{\varphi}_n \left(f_n^2 \left(\frac{c_{n+1}}{2} \xi, T \right) \right), \quad (4.17)$$

where $\tilde{f}(\xi) = \int e^{i\xi x} f(x) dx$. These facts are explained in paper [BM3]. Also the proof of Proposition 4.2 is a natural adaptation of the proof of the corresponding result (of Proposition 3) in paper [BM3]. Hence we only explain the main points and the necessary modifications.

First we remark that in the regularization of the functions $f_n(x, T)$ the same normalization could be applied as in paper [BM3]. Because of the inductive property $I(n)$ $f_n(x, T)$ is essentially concentrated in a neighbourhood of the origin of size $\sqrt{\beta_n(T)}$, and if (n, T) is in the low temperature domain and $\eta > 0$ is chosen sufficiently small, then $\frac{|x|}{100\sqrt{c^{(n)}}} \leq \frac{\eta}{10}$ for $|x| \leq \sqrt{\beta_n(T)}$, and the function $f_n(x, T)$ (disregarding the scaling with the numbers A_n and B_n) is not changing in the typical region by the regularization of the function $f_n(x, T)$. This is the reason why such a regularization works well.

The proof of Proposition 4.2 contains several estimates. First we list those results whose proof apply the bound on $f_n(x, T)$ formulated in the Inductive hypothesis $I(n)$. One can bound the differences

$$\frac{\partial^j}{\partial x^j} (\bar{\mathbf{Q}}_{n, M_n(T)}^c f_n(x, T) - \bar{\mathbf{Q}}_{n, M_n(T)}^c \varphi_n(f_n(x, T))) \quad (\text{Lemma 4 in [BM3]}),$$

$$\frac{\partial^j}{\partial x^j} (\bar{\mathbf{Q}}_{n, M_n(T)}^c \varphi_n(f_n(x, T)) - \bar{\mathbf{T}}_{n, M_n(T)}^c \varphi_n(f_n(x, T))) \quad (\text{Lemma 5 in [BM3]}),$$

with the help of Property $I(n)$ similarly to paper [BM3]. The absolute value of these expressions can be bounded for all $\varepsilon > 0$ by

$$\frac{\beta_n(T)}{c^{(n)}} \frac{C_1(\varepsilon)C^2}{\beta_n^{(j+1)/2}(T)} \exp \left\{ -\frac{2(1-\varepsilon)}{c_{n+1}\sqrt{\beta_n(T)}} \left| 2x + \frac{x^2}{c^{(n+1)}M_n(T)} \right| \right\}$$

with some appropriate constant $C_1(\varepsilon) > 0$ if $f_n(x, T)$ satisfies Condition $I(n)$.

The main difference between these estimates and the analogous results in paper [BM3] is that the upper bounds given for the above expressions contain a small multiplying factor $\frac{\beta_n(T)}{c^{(n)}}$. In paper [BM3] the multiplying factors 2^{-n} and $1/c^{(n)}$ appear instead of this term. In the proof of this paper we had to make some modifications, because while

in paper [BM3] only very low temperatures were considered when $M_n(T)$ is strongly separated from zero, now we want to give an upper bound under the weaker condition formulated in the definition of the low temperature region. The proofs are very similar. The only essential difference is that in the present case the typical region, where a good asymptotic approximation must be given is chosen as the interval $|x| < 10\sqrt{c^{(n)}}$, i.e. it does not depend on the value of $M_n(T)$.

Also the expression $\bar{\mathbf{Q}}_{n,M_n(T)}^{\mathbf{c}} f_n(x, T)$ can be bounded together with their first two derivatives with the help of Property $I(n)$ in the same way as in Lemma 3 of paper [BM3]. But this estimate is useful only for large x . It can be proved, similarly to the proof of the corresponding result in paper [BM3] (lemma 7) that the scaling constants which appear in the formulas expressing $\mathbf{Q}_{n,M_n(T)}^{\mathbf{c}}$ through $\bar{\mathbf{Q}}_{n,M_n(T)}^{\mathbf{c}}$ and $\mathbf{T}_n$ through $\bar{\mathbf{T}}_{n,M_n(T)}^{\mathbf{c}}$ are very close to each other. Here again the multiplying factor $\frac{\beta_n(T)}{c^{(n)}}$ appears in the error term instead of the multiplying factor $1/c^{(n)}$ in paper [BM3]. This Lemma 7 in [BM3] is a technical result which expresses the difference of the functions $\bar{\mathbf{T}}_{n,M_n(T)}^{\mathbf{c}} F_1(x)$ and $\bar{\mathbf{T}}_{n,M_n(T)}^{\mathbf{c}} F_2(x)$ together with its derivatives if we have a control on the difference of the original functions $F_1(x)$ and $F_2(x)$. We gain such kind of information from the inductive hypothesis $I(n)$. They give a good control on the difference $f_{n+1}(x, T) - \mathbf{T}_n \varphi_n(f_n(x, T))$. The consequences of these results are formulated in Proposition 2 in paper [BM3]. These results also imply an estimate on the Fourier transforms $\tilde{\varphi}_{n+1}(f_{n+1}(\xi, T)) - \tilde{\mathbf{T}}_n \tilde{\varphi}_n(f_n(\xi, T))$ and $\tilde{\mathbf{T}}_n \tilde{\varphi}_n(f_n(\xi, T))$ and also on their analytic continuation. This is done in lemma 8 in paper [BM3]. Now again the analogous result holds under the conditions of the present paper with the difference that the term c^{-n} must be replaced $\frac{\beta_n(T)}{c^{(n)}}$. The estimate obtained for $\tilde{\mathbf{T}}_n \tilde{\varphi}_n(f_n(\xi, T))$ in such a way is relatively weak, it is useful only for large ξ .

The above results are not sufficient to prove Proposition 4.2. In particular, they do not explain why the right scaling was chosen in the definition of the function $f_n(x, T)$. Their role is to bound the error which is committed when $\mathbf{Q}_{n,M_n(T)}^{\mathbf{c}} f_n(x, T)$ is replaced by $\mathbf{T}_n \varphi(f_n(x, T))$. The function $\mathbf{T}_n \varphi(f_n(x, T))$ together with its derivatives and Fourier transform can be well bounded by means of formula (4.17) and the inverse Fourier transform. In the estimations leading to such bounds the inductive hypothesis $J(n)$ plays a crucial role. The proof of Lemma 9 in paper [BM3] can be adapted to the present case without any essential difficulty. But, the parameters α_n , β_n and c must be replaced by $\alpha_n(T)$, $\beta_n(T)$ and c_{n+1} in the present case.

Proposition 4.2 can be proved similarly to its analog, Proposition 3 in paper [BM3]. The notation must be adapted to the notation of the present paper. Beside this, the small coefficient $c^{-n/2}$ appearing in the proof of Proposition 3 in [BM3] must be replaced by $\sqrt{\frac{\beta_n(T)}{c^{(n)}}}$. There is one point where a really new argument is needed in the proof. This argument requires a more detailed discussion. It is the proof of relation (4.14), i.e. of the fact that $\alpha_n(T)$ and $\beta_n(T)$ have the same order of magnitude. Their ratio must be bounded by a number independent of η . The proof of the analogous result

in paper [BM3] exploited the fact that in the model of that paper the sequence $c^{(n)}$ tended to infinity exponentially fast. In the present case this property does not hold any longer, hence a different argument is needed. The validity of relation (4.14) has a different cause for relatively small and large indices n .

For large n it can be shown that both $\beta_n(T)$ and $\alpha_n(T)$ have the same order of magnitude as $M_n^{-2}(T)$, and for large n these relations imply (4.14). If n is relatively small and $M_0(T)$ is large, then $M_n^{-2}(T)$ is much less than $\alpha_n(T)$ and $\beta_n(T)$. In this case the above indicated argument does not work, but it can be proved that for such indices n the numbers $\beta_n(T)$ are decreasing exponentially fast, and the proof of relation (4.14) for such n is based on this fact.

To distinguish between small and large indices n define the number

$$N_1(T) = \left\{ \min n : n \geq N, \text{ and } \beta_{n+1}(T) \leq \frac{100}{M_n^2(T)} \right\}, \quad (4.18)$$

($N_1(T) = \infty$ if there is no such n).

where the number N was defined in formula (1.22). We shall later see that $N_1(T) < \infty$ for all $0 < T \leq c_0 A_0/2$.

First we prove relation (4.14) under the additional condition $n \leq N_1(T)$. In this case $\beta_{m+1}(T) \leq \frac{c_{m+1}^2}{2} \beta_m(T) + \frac{\beta_m(T)}{10}$ for $m \leq n$, and because of Condition 1

$$\beta_{m+1}(T) \leq \frac{2}{3} \beta_m(T) \quad \text{if } m \leq N_1(T) \quad (4.19)$$

for all $N \leq m \leq n$. Hence $\sqrt{\frac{\beta_{m+1}(T)}{c^{(m+1)}}} \leq \frac{5}{6} \sqrt{\frac{\beta_m(T)}{c^{(m)}}}$, $\sqrt{\frac{\beta_m(T)}{c^{(m)}}} \leq \left(\frac{5}{6}\right)^{m-N} \sqrt{\frac{\beta_N(T)}{c^{(N)}}}$,

$$\begin{aligned} 1 \leq \frac{\beta_{m+1}(T)}{\alpha_{m+1}(T)} &\leq \max \left(\frac{\frac{c_{m+1}^2}{2} + \sqrt{\frac{\beta_m(T)}{c^{(m)}}}}{\frac{c_{m+1}^2}{2} - \sqrt{\frac{\beta_m(T)}{c^{(m)}}}} \cdot \frac{\beta_m(T)}{\alpha_m(T)}, 10^{13} \right) \\ &\leq \max \left(\exp \left\{ 5 \sqrt{\frac{\beta_m(T)}{c^{(m)}}} \right\} \cdot \frac{\beta_m(T)}{\alpha_m(T)}, 10^{13} \right). \end{aligned}$$

for $N \leq m \leq n$, and

$$\frac{\beta_{n+1}(T)}{\alpha_{n+1}(T)} \leq \max \left(\frac{\beta_N(T)}{\alpha_N(T)}, 10^{13} \right) \exp \left\{ 5 \sum_{m=N}^n \sqrt{\frac{\beta_m(T)}{c^m(T)}} \right\} \leq K.$$

The above argument together with the observation that $\beta_N(T) \gg M_N^{-2}(T)$ if the parameter $t > 0$ in (1.4) is sufficiently small and $T \leq c_0 A_0/2$ imply that $N < N_1(T)$, and the pair (n, T) is in the low temperature region for all $n \leq N_1(T)$. The latter property

follows from the fact that by formula (4.19) the sequence $\frac{\beta_n(T)}{c^{(n)}}$ is monotone decreasing for $N \leq n \leq N_1(T)$.

In the case $n > N_1(T)$ we can prove by induction with respect to n together with the inductive proof of Proposition 4.2 that

$$\beta_{n+1}(T) \leq \frac{100}{M_n^2(T)} \quad \text{if } n \geq N_1(T) \text{ and } (n, T) \text{ is in the low temperature region.} \quad (4.20)$$

By applying formula (4.20) for $n-1$ and the fact that (n, T) is in the low temperature region we get that the term $\gamma_{n-1}(T)$ in formula (4.13) can be bounded as

$$|\gamma_{n-1}(T)| \leq \frac{\beta_n(T)}{c^{(n)}} \sqrt{\beta_n(T)} \leq \eta \frac{10}{M_{n-1}(T)} \leq \frac{1}{8C_1 M_{n-1}(T)} \quad (4.21)$$

with the same number C_1 which appears in (4.13) if the number $\eta > 0$ was chosen sufficiently small. Then formula (4.13) implies that $M_n(T) \leq M_{n+1}(T)$. Hence we get by applying again formula (4.20) with $n-1$ that $M_n(T) < M_{n-1}(T)$, and

$$\beta_{n+1}(T) \leq \frac{2}{3}\beta_n(T) + \frac{10}{M_n^2(T)} \leq \frac{200}{3M_{n-1}^2(T)} + \frac{10}{M_n^2(T)} \leq \frac{100}{M_n^2(T)}.$$

This means that formula (4.20) also holds for n . Relation (4.20) together with the definition of the sequence $\alpha_n(T)$ implies that for $n \geq N_1(T)$

$$\alpha_{n+1}(T) \geq \frac{10^{-12}}{M_n^2(T)} \geq 10^{-14}\beta_n(T),$$

i.e. formula (4.14) is also valid for $n > N_1(T)$ if (n, T) is in the low temperature domain. With the help of this argument Proposition 4.2 can be proved by an adaptation of the proof of the corresponding result in [BM3].

We formulate and prove a lemma which describes some properties of the numbers $\beta_n(T)$ in the cases when $n \leq N_1(T)$ or $n \geq N_1(T)$. Several parts of it were already proved in the previous arguments.

Lemma 4.3. *Let $0 < T \leq c_0 A_0/2$. If the parameter $\kappa > 0$ in formula (1.4) is sufficiently small, then the number $N_1(T)$ defined in (4.18) is finite, and $N_1(T) > N$. The pair $(N_1(T), T)$ is in the low temperature domain. The relations (4.19), (4.20),*

$$M_n(T) - \frac{3}{8c^{(n)}M_n(T)} \leq M_{n+1}(T) \leq M_n(T) - \frac{1}{8c^{(n)}M_n(T)} \quad (4.22)$$

if $n \geq N_1(T)$ and (n, T) is in the low temperature region,

Also the relations

$$\begin{aligned} M_n(T) - \frac{1}{4c^{(n)}M_n(T)} - \eta \left(\frac{2}{3}\right)^{(n-N)/2} &\leq M_{n+1}(T) \\ &\leq M_n(T) - \frac{1}{4c^{(n)}M_n(T)} + \eta \left(\frac{2}{3}\right)^{(n-N)/2} \quad \text{if } N \leq n \leq N_1(T), \end{aligned} \quad (4.23)$$

and

$$N_1(T) - N \leq 10 \log(1/\kappa T^2) \quad \text{with the parameter } \kappa \text{ appearing in (1.17)}. \quad (4.24)$$

hold. If $M_n(T) < 10$ then $n \geq N_1(T)$.

Proof. Formulas (4.19) and (4.20) were already proved in the previous argument, and since (N, T) is in the low temperature region, i.e. $\beta_N(T) \geq \eta c^N$, relation (4.19) implies that (n, T) is in the low temperature region for all $N \leq n \leq N_1(T)$. Formula (4.22) follows from formula (4.21) with the replacement of $n - 1$ by n and formula (4.13). By relation (4.19) $\beta_n(T) \leq \left(\frac{2}{3}\right)^{n-N}$ if $N \leq n \leq N_1(T)$. Hence it follows from (4.13) that

$$M_{n+1}(T) \leq M_n(T) + \frac{\beta_{n+1}(T)}{c^{(n)}} \frac{\sqrt{\beta_{n+1}(T)}}{c^{(n)}} \leq M_n(T) + \eta \left(\frac{2}{3}\right)^{(n-N)/2}, \quad (4.25)$$

and even relation (4.23) holds in this case.

Relation (4.25) and the estimate obtained for $\beta_n(T)$ imply that $M_n^2(T) \leq (M_N(T) + 1)^2 \leq 2M_N^2(T)$ and $\beta_{n+1}(T)M_n^2(T) \leq 2M_N^2(T) \left(\frac{2}{3}\right)^{n-N}$ if $n \leq N_1(T)$. This relation together with the definition of the index $N_1(T)$ defined in (4.18) imply that $2M_N^2(T) \left(\frac{2}{3}\right)^{n-N} \geq 100$ if $n < N_1(T)$. Applying the last formula for $n = N_1(T) - 1$ we get that $(N_1(T) - 1 - N) \log \frac{3}{2} \leq \log \frac{M_N^2(T)}{50}$. Since $M_N^2(T) \sim \text{const.} \frac{1}{\kappa T^2}$ this relation implies that $N_1(T)$ is finite, and moreover it satisfies (4.24). Finally, if the inequalities $M_n(T) \leq 10$ and $n < N_1(T)$ held simultaneously, then the inequality $M_n^2(T)\beta_{n+1}(T) \leq 100 \left(\frac{2}{3}\right)^{n-N} \leq 100$ would also hold. This relation contradicts to the assumption $n < N_1(T)$. Hence also the last statement of Lemma 4.3 holds.

The previous results enable us to describe the different behaviour of the model in the cases when the Dyson condition (1.3) is satisfied and when it is not. This will be done in Lemma 4.4. It shows that if (1.3) *is not satisfied* then for all T there is a pair (n, T) which does not belong to the low temperature region, while if (1.3) *is satisfied*, then all sufficiently low temperatures T belong to the low temperature region. In the latter case the asymptotic behaviour of the spontaneous magnetization $M_n(T)$ can be described for large n . The description of the behaviour of the function $q_n(x, T)$ in the case when T does not belong to the low temperature region needs further investigation, and this will be done in Sections 5 and 6. A more detailed investigation of the case when T belongs to the low temperature region will be done in Section 7. We finish this section with the proof of a result about the behaviour of the magnetization $M_n(T)$ at low temperatures $T > 0$ which will be useful in the subsequent part of the paper.

Lemma 4.4. *Let $0 < T \leq c_0 A_0/2$, and let the parameter $\kappa > 0$ in formula (1.17) be sufficiently small. If the Dyson condition (1.3) is not satisfied, then for all $T > 0$ there is some $n = n(T)$ for which (n, T) does not belong to the low temperature region. If, on the other hand, condition (1.3) is satisfied, then T belongs to the low temperature region for sufficiently small $T > 0$. In this case relation (3.8) and under the additional Condition 2 also relation (3.9) (with $r = 2$) hold.*

Proof. It follows from formulas (4.22) and (4.23) that

$$-\frac{1}{c^{(n)}} \leq M_{n+1}^2(T) - M_n^2(T) \leq -\frac{1}{8c^{(n)}} \quad (4.26)$$

if $n \geq N_1(T)$ and the pair (n, T) is in the low temperature region, and

$$\begin{aligned} -\frac{1}{2c^{(n)}} - 10 \left(\frac{2}{3}\right)^{n-N} (M_N(T) + 1) &\leq M_{n+1}^2(T) - M_n^2(T) \\ &\leq -\frac{1}{2c^{(n)}} + 10 \left(\frac{2}{3}\right)^{n-N} (M_N(T) + 1) \end{aligned} \quad (4.26')$$

if $N \leq n \leq N_1(T)$. Formula (4.26) can be obtained by taking square in formula (4.22) and observing that $c^{(n)} M_n(T)^2 > 10\eta^{-1}$. Formula (4.26') can be deduced similarly from (4.23) by observing first that the right-hand side of (4.23) implies that $M_n(T) \leq M_N(T) + 1$ for $N \leq n \leq N_1(T)$.

Formulas (4.26) and (4.26') imply that if a temperature $T > 0$ is in the low temperature region, then

$$\sum_{k=N}^n \frac{1}{c^{(k)}} \leq 8(M_N^2(T) - M_n^2(T)) + 30(M_N(T) + 1) \leq 8M_N^2(T) + 30(M_N(T) + 1)$$

for all $n \geq N$, where the number N is defined in (1.22). Since the right-hand side of the last formula does not depend on n , this implies that (1.3) holds.

In the other direction, if (1.3) holds, then since by Proposition 4.1 $\lim_{T \rightarrow \infty} M_0(T) = \lim_{T \rightarrow \infty} M_N(T) = \infty$, there is some number $\bar{T} \leq c_0 A_0/2$ such that for all temperatures $0 < T \leq \bar{T}$ $M_N^2(T) > 8 \sum_{n=N}^{\infty} \frac{1}{c^{(n)}} + 30M_N(T) + 31$. If $T > 0$ satisfies the above inequality, then the left-hand side of the inequalities (4.26) and (4.26') imply that if the pair (n, T) is in the low temperature domain and $n \geq N_1(T)$, then

$$M_n^2(T) > M_N^2(T) - 8 \sum_{n=N}^n \frac{1}{c^{(n)}} 30(M_N(T) + 1) \geq 1.$$

Hence $M_n^2(T) > 1$ for all n , and T is in the low temperature region.

Let $T > 0$ be in the low temperature region. If $n > m > N_1(T)$, then by (4.26)

$$|M_n^2(T) - M_m^2(T)| \leq \sum_{k=m}^n \frac{1}{c^{(k)}}.$$

Since in this case Condition 1 holds, the last relation implies that $M_n^2(T)$, $n = 1, 2, \dots$, is a Cauchy sequence, and relation (3.8) holds. We claim that if Condition 2 also holds, then for any $\varepsilon > 0$

$$-\frac{1+\varepsilon}{2c^{(n)}} \leq M_{n+1}^2(T) - M_n^2(T) \leq -\frac{1-\varepsilon}{2c^{(n)}} \quad (4.27)$$

if $n \geq n(\varepsilon)$. Relation (3.9) is a consequence of (4.27). Relation (4.27) can be deduced from (4.13) and (4.20) if we show that for any temperature $T > 0$ in the low temperature region

$$\lim_{n \rightarrow \infty} \frac{\beta_n(T)}{c^{(n)}} = 0. \quad (4.28)$$

Relation (4.28) holds under Condition 2, since by (4.26) in this case for all $n > N_1(T)$

$$M_n^2(T) \geq \lim_{k \rightarrow \infty} (M_n^2(T) - M_k^2(T)) \geq \frac{1}{8} \sum_{k=n}^{\infty} \frac{1}{c^{(k)}},$$

and $\frac{\beta_n(T)}{c^{(n)}} \leq \frac{100}{M_{n-1}^2(T)c^{(n)}} \leq 800 \left(c^{(n)} \sum_{k=n-1}^{\infty} \frac{1}{c^{(k)}} \right)^{-1}$. Under Condition 2 the last expression tends to zero as $n \rightarrow \infty$. This implies formula (4.27). Lemma 4.4 is proved.

5. Estimates in the Intermediate Region. Proof of Theorem 3.1

In this section we give some estimates on $q_n(x, T)$ when the pair (n, T) belongs neither to the low nor to the high temperature region and prove Theorem 3.1 with their help.

Let us consider the number $\bar{n} = \bar{n}(T)$ introduced in the formulation of Theorem 3.1. We shall prove some estimates about a scaled version of the function $q_{\bar{n}(T)}(x, T)$ in Lemmas 5.1 and 5.2. In Lemma 5.1 the case $T \leq c_0 A_0$, in Lemma 5.2 the case $T \geq c_0 A_0$ will be considered. Lemmas 5.1 and 5.2 yield some estimates on the tail-behaviour of a scaled version of the function $q_{\bar{n}(T)}(\cdot, T)$. This will be needed to start an inductive procedure for all $n \geq \bar{n}(T)$ which state that the functions $q_n(x, T)$ become more and more strongly concentrated around zero as the index n is increasing. This procedure is based on Lemmas 5.3 and 5.4. The role of Lemma 5.3 is to give an appropriate lower bound for the norming constant $Z_n(T)$ in the definition of the function $q_n(x, T)$. Then in Lemma 5.4 we prove some contraction property of the operator which maps an appropriate scaled version of the distribution function with density function $\text{const. } q_{n-1}(|x|, T)$ to an appropriate scaled version of the distribution function with density $\text{const. } q_n(|x|, T)$, $x \in \mathbb{R}^2$. The proof of Lemma 5.4 will exploit the rotation symmetry of the model. Theorem 3.1 will be proved by means of these lemmas.

To formulate these results we introduce some notations.

Let us introduce the functions

$$\hat{h}_n(x, T) = \frac{1}{\sqrt{c^{(\bar{n}(T))}}} q_n \left(\frac{x}{\sqrt{c^{(\bar{n}(T))}}}, T \right), \quad x \in \mathbb{R}^2 \quad (5.1)$$

and measures

$$\hat{H}_{n,T}(\mathbf{A}) = \int_{\mathbf{A}} \hat{h}_n(x, T) dx, \quad \mathbf{A} \subset \mathbb{R}^2 \quad (5.2)$$

in the space $\mathbb{R}^2$. Define also the function

$$\hat{H}_{n,T}(R) = \hat{H}_{n,T}(\{x: |x| \geq R\}) \quad \text{for } R \geq 0. \quad (5.3)$$

The functions $\hat{h}_{n,T}$ and measures $\hat{H}_{n,T}$ are similar to the functions $h_{n,T}$ and measures $H_{n,T}$ defined in (3.5) and (3.6). The only difference is that the scaling of $q_n(x, T)$ in (5.2) and (5.3) is made by means of $c^{(\bar{n}(T))}$ instead of $c^{(n)}$. If Condition 5 is satisfied with a sufficiently small $\bar{\eta}$ and sufficiently large $L(\bar{\eta}, t)$, and $n - \bar{n}(T)$ is not too large, then the approximation of $c^{(n)}$ by $c^{(\bar{n}(T))}$ is sufficiently good for our purposes. Hence it will be enough to have a good control on the measure $\hat{H}_{n,T}$. In Lemma 5.3 we give a bound on it for large $|x|$ and in Lemma 5.4 we prove an estimate which enables to bound $\hat{H}_{n,T}(x)$ for small x too.

With the help of these results we can prove that starting from $\bar{n} = \bar{n}(T)$ after finitely many steps k the pair $(\bar{n} + k, T)$ is in the high temperature region. Moreover, this number k can be bounded from above independently of the temperature T . First we formulate Lemma 5.1.

Lemma 5.1. *Under the conditions of Proposition 4.2 the function $h_{\bar{n}(T)}(x, T)$ defined in (3.5) satisfies the inequality*

$$h_{\bar{n}(T)}(x, T) \leq \exp \left\{ \frac{K}{\eta} - \frac{|x|^2}{10} \right\} \quad \text{if } T \leq c_0 A_0 / 2 \quad (5.4)$$

with an appropriate $K > 0$. For $T \leq c_0 A_0 / 2$ the pair $(\bar{n}(T), T)$ does not belong to the high temperature region, and there exists some $\tilde{\eta} = \tilde{\eta}(\eta)$ such that the function $\hat{H}_{\bar{n}, T}(\cdot)$ defined in (5.3) satisfies the inequality

$$\hat{H}_{\bar{n}(T), T}(\tilde{\eta}^{-1}) \leq 1/2 \quad \text{if } T \leq c_0 A_0 / 2, \quad (5.5)$$

i.e. for $T \leq c_0 A_0 / 2$ there is a circle with its center in the origin whose radius depends only on η , and whose $\hat{H}_{\bar{n}(T), T}$ measure is greater than $1/2$.

Proof. Let us introduce the function

$$\bar{h}_n(x, T) = \frac{1}{\sqrt{c^{(n)}}} \bar{q}_n \left(\frac{x}{\sqrt{c^{(n)}}}, T \right), \quad x \geq 0$$

with the function $\bar{q}_n$ introduced in (2.17). This function is very similar to the intersection of the function $h_n(x, T)$ with the coordinate axis $y = 0$. Only the norming of the two functions is different, since $\int_{\mathbb{R}^2} h_n(x, T) dx = 1$, and $\int_0^\infty \bar{h}_n(x, T) dx = 1$.

We can apply Proposition 4.2 with the choice $n = \bar{n}(T) - 1$. Since hypothesis $I(n)$ holds for $n = \bar{n}$

$$f_{\bar{n}(T)}(x, T) \leq \frac{K}{\beta_{\bar{n}(T)-1}^{1/2}(T)} \exp \left\{ -\frac{1}{\sqrt{\beta_{\bar{n}(T)}(T)}} \left| 2x + \frac{x^2}{c^{(\bar{n}(T))} M_{\bar{n}(T)}(T)} \right| \right\} \\ \text{if } x > -c^{(\bar{n}(T))} M_{\bar{n}(T)}(T)$$

with some universal constant $K > 0$. It follows from this relation that the function $\bar{h}_{\bar{n}(T)}(x, T) = \sqrt{c^{(\bar{n}(T))}} f_{\bar{n}(T)} \left(\sqrt{c^{(\bar{n}(T))}} x - c^{(\bar{n}(T))} M_{\bar{n}(T)}(T), T \right)$ satisfies the inequality

$$\bar{h}_{\bar{n}(T)}(x, T) \leq K \left(\frac{c^{(\bar{n}(T))}}{\beta_{\bar{n}(T)-1}(T)} \right)^{1/2} \exp \left\{ \frac{1}{\sqrt{\beta_{\bar{n}(T)}(T)}} \left(c^{(\bar{n}(T))} M_{\bar{n}(T)}(T) - \frac{x^2}{M_{\bar{n}(T)}(T)} \right) \right\}$$

The inequalities $\beta_{\bar{n}(T)}(T) > \eta c^{(\bar{n}(T))}$ and $\beta_{\bar{n}(T)-1}(T) \leq \eta c^{(\bar{n}(T)-1)}$ hold. Lemma 4.3 implies that the fractions $\frac{\beta_{\bar{n}(T)}(T)}{\beta_{\bar{n}(T)-1}(T)}$, $\frac{M_{\bar{n}(T)}(T)}{M_{\bar{n}(T)-1}(T)}$ and $\beta_{\bar{n}(T)}(T) M_{\bar{n}(T)}(T)^2$ are separated

both from zero and infinity. Hence $\frac{c^{(\bar{n}(T))}}{\beta_{\bar{n}(T)-1}(T)} \leq \frac{\text{const.}}{\eta}$, $\frac{c^{(\bar{n}(T))} M_{\bar{n}(T)}(T)}{\sqrt{\beta_{\bar{n}(T)}(T)}} \leq \frac{\text{const.}}{\eta}$

and $\frac{1}{M_{\bar{n}(T)}(T)\sqrt{\beta_{\bar{n}(T)}(T)}} \geq \frac{1}{20}$. These inequalities together with the last relation imply that

$$\bar{h}_{\bar{n}(T)}(x, T) \leq e^{\bar{K}/\eta} e^{-x^2/20} \quad (5.6)$$

with an appropriate $\bar{K} > 0$. Since the relation

$$h_{\bar{n}(T)}(x, T) = C(T)\bar{h}_{\bar{n}(T)}(x, T) \quad (5.7)$$

holds between the functions $h_{\bar{n}(T)}$ and $\bar{h}_{\bar{n}(T)}$ with an appropriate number $C(T)$, formula (5.4) can be deduced from (5.6) if we give a good upper bound for the constant $C(T)$ in (5.7). Observe that

$$\int_{\mathbb{R}^2} \bar{h}_{\bar{n}(T)}(|x|, T) dx = 2\pi \int_0^\infty x \bar{h}_{\bar{n}(T)}(x, T) dx \geq 2\pi R \left(1 - \int_0^R \bar{h}_{\bar{n}(T)}(x, T) dx\right)$$

for any $R > 0$, and by formula (5.6)

$$\int_0^R \bar{h}_{\bar{n}(T)}(x, T) dx \leq \frac{1}{2}$$

if $0 < T \leq c_0 A_0/2$ and $R = \frac{1}{2}e^{-K/\eta}$. Hence $C(T)^{-1} = \int_{\mathbb{R}^2} h_{\bar{n}(T)}(x, T) dx \geq e^{-K/\eta}$. This means that $C(T) \leq e^{K/\eta}$ in (5.7), and inequality (5.6) follows from (5.4), only the constant K in (5.6) must be replaced by $2K$. We also need a lower bound for $C(T)$ in (5.7). To get it observe that

$$\int_{\mathbb{R}^2} \bar{h}_{\bar{n}(T)}(|x|, T) dx = 2\pi \int_0^\infty x \bar{h}_{\bar{n}(T)}(x, T) dx = M_{\bar{n}(T)} \sqrt{c(\bar{n}(T))} \leq \frac{10}{\sqrt{\eta}}.$$

This inequality implies that $C(T) \geq \frac{\sqrt{\eta}}{10}$ in (5.7) and

$$\begin{aligned} D_{\bar{n}(T)}^2(T) &= \int_{\mathbb{R}^2} |x|^2 h_{\bar{n}(T)}(x, T) dx = 2\pi \int_0^\infty x^3 h_{\bar{n}(T)}(x, T) dx \\ &\geq 2\pi \frac{\sqrt{\eta}}{10} \int_0^\infty x^3 \bar{h}_{\bar{n}(T)}(|x|, T) dx \geq 2\pi \frac{\sqrt{\eta}}{10} \left(\int_0^\infty x \bar{h}_{\bar{n}(T)}(|x|, T) dx \right)^3 \geq \frac{\text{const.}}{\eta}. \end{aligned}$$

This implies that the pair $(\bar{n}(T), T)$ is not in the high temperature region. Finally, it follows from (5.4) that $H_n(R) \leq 1/2$ for $R = e^{2K/\eta}$. Lemma 5.1 is proved.

If $T \geq c_0 A_0/2$, then $\bar{n}(T) = 0$, and $\bar{h}_{\bar{n}(T)}(x, T) = c_0^{-1/2} \bar{q}_0(c_0^{-1/2} x, T)$, where $\bar{q}_0(x, T)$ is defined in (2.13) and (2.14). Hence

$$\bar{h}_{\bar{n}(T)}(x, T) = \frac{1}{Z_0(T)} \exp \left\{ \left(A_0 - \frac{T}{\bar{c}_0} \right) \frac{x^2}{2} - \kappa T^2 \frac{x^4}{4c_0^2} \right\} \quad \text{if } T \geq c_0 A_0/2 \quad (5.8)$$

with the norming constant

$$Z_0(T) = 2\pi \int_0^\infty x \exp \left\{ \left(A_0 - \frac{T}{\bar{c}_0} \right) \frac{x^2}{2} - \kappa T^2 \frac{x^4}{4c_0^2} \right\} dx. \quad (5.8')$$

With the help of formulas (5.8) and (5.8') we shall prove the following

Lemma 5.2. *There exists $\kappa_0 = \kappa_0(N) > 0$ such that if $0 < \kappa < \kappa_0$ and $T \geq c_0 A_0/2$, then $\bar{n}(T) = 0$, and*

$$h_{\bar{n}(T)}(x, T) \leq 10T \exp \left\{ -10Tx^2 + \frac{100}{\kappa} \right\} \quad \text{if } T \geq c_0 A_0/2 \quad (5.9)$$

$$h_{\bar{n}(T)}(x, T) \leq 10T \quad \text{if } T \geq c_0 A_0/2 \quad (5.9')$$

$$h_{\bar{n}(T)}(x, T) \leq 100e^{-Tx^2/4} \quad \text{if } T \geq 10A_0 \text{ and } |x| \geq T^{-1/3}. \quad (5.9'')$$

The pair $(\bar{n}(T), T)$ belongs to the high temperature region if T is very large, e.g. if $T \geq e^{-1/\eta^9}$, and it does not belong to it if $T > 0$ is relatively small, e.g. if $T \leq \eta^{-100}$. If $(\bar{n}(T), T)$ does not belong to the high temperature region, then the function $h_{\bar{n}(T)}(x, T)$ defined in formula (3.5) satisfies the inequality

$$h_{\bar{n}(T)}(x, T) \leq \exp\{K(\eta, \kappa) - \alpha|x|^2\} \quad (5.10)$$

with a constant $\alpha = \alpha(\eta) > 0$ and an appropriate number $K(\eta, \kappa)$ depending only on κ and η . In this case there is a constant $B = B(\eta, \kappa) > 0$ in such a way that the quantity $\hat{H}_{\bar{n}(T), T}(\cdot)$ defined in (5.3) satisfies the inequality

$$1 - \hat{H}_{\bar{n}(T), T}(B) \leq \frac{1}{2}. \quad (5.11)$$

This means that if the pair $(\bar{n}(T), T)$ is not in the high temperature region (and $T \geq c_0 A_0/2$), then there is a radius $B = B(\eta, \kappa)$ such that the $\hat{H}_{\bar{n}(T), T}$ measure of the circle $\{x: |x| \leq B(\eta, \kappa)\}$ which is bigger than half.

If $(\bar{n}(T), T) = (0, T)$ is in the high temperature region, then

$$\hat{H}_{\bar{n}(T), T}(x) \leq K_1 e^{-K_2 \eta^2 x^2} \quad \text{for all } x > 0 \quad (5.12)$$

with some universal constants $K_1 > 0$ and $K_2 > 0$.

Proof. First we estimate the norming factor $Z_0(T)$ from below. Let us observe that $\left(A_0 - \frac{T}{c_0}\right) \frac{x^2}{2} - \kappa T^2 \frac{x^4}{4c_0^2} \geq -10Tx^2$ if $\kappa T x^2 \leq 1/100$ and $c_0 A_0/2 \leq T$. Hence

$$Z_0(T) \geq 2\pi \int_0^{1/10\sqrt{\kappa T}} x e^{-10Tx^2} dx = 2\pi \int_0^{1/10\sqrt{\kappa}} \frac{x e^{-10x^2}}{T} dx \geq \frac{1}{10T}. \quad (5.13)$$

If $T \geq c_0 A_0/2$, then $\left(A_0 - \frac{T}{c_0}\right) \frac{x^2}{2} - \kappa T^2 \frac{x^4}{4c_0^2} \leq \frac{100}{\kappa} - 10Tx^2$. This relation together with (5.13) imply formula (5.9), and formula (5.9') follows from (5.13).

If $T \geq 10A_0$ then $\left(A_0 - \frac{T}{c_0}\right) \frac{x^2}{2} - \kappa T^2 \frac{x^4}{4c_0^2} \leq -\frac{T}{2}x^2 \leq -\frac{T}{4}x^2 - \frac{T^{1/3}}{4}$ for $|x| \geq T^{-1/3}$.

This relation together with (5.13) imply relation (5.9'').

Formula (5.9'') implies that if $T > e^{-1/\eta^9}$, then the pair $(0, T)$ belongs to the high temperature region. To see that for $T < \eta^{-100}$ the pair $(0, T)$ does not belong to the high temperature domain it is enough to observe that in this case by formula (5.9') the $H_{0,T}$ measure of the circle $\{x: |x| \leq \eta^{100}\}$ is less than $10\pi T \eta^{200} \leq 1/2$. Hence in this case the variance $D_0^2(T)$ is larger than in the high temperature region. Inequality (5.9) together with the fact that if the pair $(0, T)$ does not belong to the high temperature region then $T \leq e^{-1/\eta^9}$ imply relations (5.10) and (5.11).

Since $T > \eta^{-100}$ if the pair $(0, T)$ is in the high temperature region, relation (5.9'') implies relation (5.12). Lemma (5.2) is proved.

To prove Lemmas 5.3 and 5.4 we rewrite formula (2.12) for the functions $\hat{h}_n(x, T)$ defined in (5.1). It has the form

$$\hat{h}_{n+1}(x, T) = \frac{2}{Z_n(T)} \int_{\mathbb{R}^2} \exp \left\{ -\frac{c^{(n)}}{c^{(\bar{n}(T))}} u^2 \right\} \hat{h}_n(x - u, T) \hat{h}_n(x + u, T) du \quad (5.14)$$

with

$$Z_n(T) = 2 \int_{\mathbb{R}^2 \times \mathbb{R}^2} \exp \left\{ -\frac{c^{(n)}}{c^{(\bar{n}(T))}} u^2 \right\} \hat{h}_n(x - u, T) \hat{h}_n(x + u, T) du dx \quad (5.14')$$

for all $n \geq \bar{n}(T)$. Let us also introduce the moment generating function of the measures $\hat{H}_{n,T}$ defined in (5.2)

$$\varphi_{n,T}(u) = \int_{\mathbb{R}^2} e^{ux} \hat{h}_{n,T}(x) dx, \quad u = (u_1, u_2) \in \mathbb{R}^2,$$

where ux denotes scalar product. By studying the properties of the moment generating function $\varphi_{n,T}(u)$ in Lemma 5.3 we give an upper bound for the function $\hat{H}_{n,T}(R)$ for large values R .

Lemma 5.3. *There exists $\kappa_0 = \kappa_0(N)$ with the number N defined in (1.22) such that for all $0 < \kappa < \kappa_0$ a number $L = L(\kappa, \bar{\eta})$ can be chosen in such a way that if Conditions 1 and 5 are satisfied, then the following relations hold. For all temperatures $T > 0$ for which the number $\bar{n}(T)$ exists, and the pair $(\bar{n}(T), T)$ does not belong to the high temperature region, the inequality*

$$\hat{H}_{\bar{n}(T)+l,T}(x) \leq e^{-2^l \alpha |x|^2/5} \quad \text{if } |x| \geq D \text{ and } 0 \leq l \leq L \quad (5.15)$$

holds with appropriate constants $\alpha > 0$ and $D > 0$. Also the norming factor $Z_n(T)$ in (5.14') can be estimated as

$$Z_{\bar{n}(T)+l}(T) \geq 2D_1 \quad \text{for } 0 \leq l \leq L \quad (5.16)$$

with some constant $D_1 > 0$. Here $\alpha = \alpha(\eta)$, and the numbers $D > 0$ and $D_1 > 0$ do not depend on the temperature T .

Proof. It follows from formulas (5.4) and (5.10) that

$$\varphi_{\bar{n}(T),T}(u) \leq \exp \left\{ K_0 + \frac{u^2}{\alpha} \right\} \quad \text{for all } u \in \mathbb{R}^2$$

with some $K_0 = K_0(\eta, \kappa) > 100$ and $\alpha = \alpha(\eta) > 0$. It can be seen by induction with respect to l that

$$\varphi_{\bar{n}(T)+l,T}(u) \leq \exp \left\{ 2^l K_l + \frac{u^2}{2^l \alpha} \right\} \quad \text{for all } 0 \leq l \leq L \text{ and } u \in \mathbb{R}^2 \quad (5.17)$$

with

$$K_l = K_{l-1} - \frac{\log \frac{Z_{\bar{n}(T)+l-1,T}}{2}}{2^l}. \quad (5.17')$$

Indeed, the function $\hat{h}_{\bar{n}(T)+l+1,T}(x)$ is increased if the kernel term $\exp \left\{ -\frac{c^{(n)}}{c^{(\bar{n}(T))}} u^2 \right\}$ is omitted from the integral in (5.14), and the integral turns into the convolution $2\hat{h}_{\bar{n}(T)+l,T} * \hat{h}_{\bar{n}(T)+l,T}(2x)$ after this change. By computing this convolution with the help of the inductive hypothesis and dividing it by $\frac{Z_{\bar{n}(T)+l+1}}{2}$ we get an upper bound for $\varphi_{\bar{n}(T)+l+1,T}(u)$. Formulas (5.17) and (5.17') follow from these calculations. We shall prove formulas (5.15) and (5.16) from these relations by induction for l together with the inductive hypothesis that

$$K_l \leq B \quad \text{for all } 0 \leq l \leq L \quad (5.18)$$

with some constants $B > 10$ depending only on κ and $\bar{\eta}$.

By applying a standard technique for the estimation of probabilities by means of moment generating functions we get with the help of formula (5.17) that the function $\hat{H}_{\bar{n}(T)+l,T}(R)$ defined in formulas (5.2) and (5.3) satisfies the inequality

$$\begin{aligned} \hat{H}_{\bar{n}(T)+l,T}(R) &\leq 4\hat{H}_{\bar{n}(T)+l,T} \left(\left\{ x = (x_1, x_2) \in \mathbb{R}^2, x_1 > \frac{R}{\sqrt{2}} \right\} \right) \\ &\leq 4 \exp \left\{ -\frac{uR}{\sqrt{2}} + 2^l K_l + \frac{u^2}{2^l \alpha} \right\} \end{aligned}$$

for all real numbers u . In particular,

$$\hat{H}_{\bar{n}(T)+l,T}(R) \leq 4 \exp \left\{ 2^l \left(K_l - \frac{R^2 \alpha}{8} \right) \right\} \quad (5.19)$$

with the choice $u = 2^{l-3/2} R\alpha$. Hence

$$\hat{H}_{\bar{n}(T)+l,T} \left(4\sqrt{\frac{B}{\alpha}} \right) \leq 4e^{-2^l B} \leq \frac{1}{2} \quad (5.20)$$

with the number $B > 0$ appearing in (5.18). Formula (5.20) implies that

$$\hat{H}_{\bar{n}(T)+l,T} \left(\left\{ x : x \in \mathbb{R}^2, |x| \leq 4\sqrt{\frac{B}{\alpha}} \right\} \right) \geq \frac{1}{2}. \quad (5.20')$$

For $z \in \mathbb{R}^2$ and $u > 0$ let $K(z, u) = \{x : x \in \mathbb{R}^2, |x - z| \leq u\}$ denote the circle with center z and radius u . Since the circle $\left\{ x : x \in \mathbb{R}^2, |x| \leq 4\sqrt{\frac{B}{\alpha}} \right\}$ can be covered by $64B(\alpha\bar{\eta})^{-1}$ circle of radius $\sqrt{\bar{\eta}}$ there is a circle $K(z, \sqrt{\bar{\eta}})$ of radius $\sqrt{\bar{\eta}}$ whose $\hat{H}_{n,T}$ measure (this measure was defined in (5.2)) is greater than $\frac{\alpha\bar{\eta}}{128B}$. Hence

$$\hat{H}_{\bar{n}(T)+l,T} \times \hat{H}_{\bar{n}(T)+l,T} (K(z, \sqrt{\bar{\eta}}) \times K(z, \sqrt{\bar{\eta}})) \geq \frac{\alpha^2 \bar{\eta}^2}{4096B^2},$$

and because of Condition 5 the expression $2Z_n(T)$ defined in (5.14) can be bounded by means of the estimation

$$\begin{aligned} \frac{Z_{\bar{n}(T)+l}(T)}{2} &\geq \int_{x+u \in K(2z, 2\sqrt{\bar{\eta}}), u \in K(z, \sqrt{\bar{\eta}})} \exp \left\{ -\frac{c^{(\bar{n}(T)+l)} (x-u)^2}{c^{\bar{n}(T)}} \right\} \\ &\quad \hat{h}_{\bar{n}(T)+l,T}(x) \hat{h}_{\bar{n}(T)+l,T}(u) dx du \\ &\geq e^{-5\bar{\eta}} \hat{H}_{\bar{n}(T)+l,T} \times \hat{H}_{\bar{n}(T)+l,T} (K(z, \sqrt{\bar{\eta}}) \times K(z, \sqrt{\bar{\eta}})) \geq e^{-1} \frac{\alpha^2 \bar{\eta}^2}{4096B^2} \\ &\geq \frac{\alpha^2 \bar{\eta}^2}{15000B^2}. \end{aligned}$$

The last relation implies (5.16) with $D_1 = \frac{\alpha^2 \bar{\eta}^2}{15000B^2}$. We get from (5.17') and the inductive hypothesis (5.18) that

$$K_l \leq (1 - 2^{-l})B,$$

if the number B is chosen as $B = \max(K_0, K^*)$, where K^* is the larger solution of the equation $x = \log \frac{15000x^2}{\alpha^2 \bar{\eta}^2}$. This implies validity of the inductive hypothesis (5.18) for l . Finally, relation (5.15) follows from (5.18) and (5.19). Lemma 5.3 is proved.

Formulas (5.14) and (5.14') can be rewritten for the function $\hat{H}_{n,T}(R)$ defined in (5.3) as

$$\begin{aligned}\hat{H}_{n+1,T}(R) &= \frac{2}{Z_n(T)} \int_{|x| \geq R} \int_{u \in \mathbb{R}^2} \exp \left\{ -\frac{c^{(n)}}{c^{(\bar{n}(T))}} u^2 \right\} \hat{h}_n(x-u, T) \hat{h}_n(x+u, T) du dx \\ &= \frac{1}{Z_n(T)} \int_{\left| \frac{x+u}{2} \right| \geq R} \int_{u \in \mathbb{R}^2} \exp \left\{ -\frac{c^{(n)}}{c^{(\bar{n}(T))}} \frac{(x-u)^2}{4} \right\} \hat{H}_{n,T}(dx) \hat{H}_{n,T}(du)\end{aligned}\quad (5.21)$$

with

$$Z_n(T) = \int_{x \in \mathbb{R}^2} \int_{u \in \mathbb{R}^2} \exp \left\{ -\frac{c^{(n)}}{c^{(\bar{n}(T))}} \frac{(x-u)^2}{4} \right\} \hat{H}_{n,T}(dx) \hat{H}_{n,T}(du). \quad (5.21')$$

for all $R \geq 0$. We apply these formulas in the proof of the following Lemma 5.4. The proof of Lemma 5.4 also exploits the rotational invariance of the measure $\hat{H}_{n,T}$.

Lemma 5.4. *Let the conditions of Lemma 5.3 hold. Then there exist some numbers $\delta = \delta(\hat{\eta}, D_1) > 0$ and $M = M(\hat{\eta}, D_1) > 0$ depending only on the numbers D_1 in formula (5.16) and $\bar{\eta}$ in Condition 5 in such a way that*

$$\begin{aligned}\hat{H}_{\bar{n}(T)+l+1,T}((1-\delta)R) &\leq \frac{1}{2} \hat{H}_{\bar{n}(T)+l+1,T}((1-\delta)R) + M \hat{H}_{\bar{n}(T)+l+1,T}(R) \\ &\text{for all } R > 0 \text{ and } 0 \leq l \leq L.\end{aligned}\quad (5.22)$$

Proof. Observe that

$$\begin{aligned}\left\{ \left| \frac{x+u}{2} \right| \geq (1-\delta)R \right\} &\subset \{|x| \geq R\} \cup \{|u| \geq R\} \\ &\cup \{|x| \geq (1-\delta)R, \arg(x, u) \leq \alpha\} \cup \{|u| \leq (1-\delta)R, \arg(x, u) \geq \alpha\}\end{aligned}$$

for all $R > 0$ and $0 < \delta < 1$ with $\alpha = 2 \arccos(1-\delta)$. Indeed, if $\left| \frac{x+u}{2} \right| \geq (1-\delta)R$, then either $|x| > R$ or $|u| > R$ or both $|x|$ and $|u|$ is less than R , but in this case either $|x| > (1-\delta)R$ or $|u| > (1-\delta)R$, and the angle between the vectors x and u must be small. On the other hand, because of the rotational invariance of the measure $\hat{H}_{n,T}$

$$\begin{aligned}\hat{H}_{\bar{n}(T)+l,T} \times \hat{H}_{\bar{n}(T)+l,T}(\{(x, y) : |x| \geq (1-\delta)R, \arg(x, u) \leq \alpha\}) \\ \leq \frac{\alpha}{\pi} \hat{H}_{\bar{n}(T)+l,T}(\{x : |x| \geq (1-\delta)R\}) = \frac{\alpha}{\pi} \hat{H}_{\bar{n}(T)+l,T}((1-\delta)R).\end{aligned}$$

The last two relations together with (5.21) and the inequality $\frac{\alpha}{\pi} \leq \sqrt{\delta}$ imply that

$$\hat{H}_{\bar{n}(T)+l+1,T}((1-\delta)R) \leq \frac{1}{Z_n(T)} \left(2\sqrt{\delta} \hat{H}_{\bar{n}(T)+l,T}((1-\delta)R) + 2\hat{H}_{\bar{n}(T)+l,T}(R) \right). \quad (5.23)$$

Relation (5.22) follows from (5.23) and (5.16) if we choose $\delta > 0$ so small that the inequality $\frac{2\sqrt{\delta}}{D_1} \leq \frac{1}{2}$ holds. Lemma 5.4 is proved.

Put $P(j, l) = P(j, l, T) = \hat{H}_{\bar{n}(T)+l}((1-\delta)^j D)$, $j = 0, 1, \dots$, $0 \leq l \leq L$ with the number D appearing in (5.15) and δ in Lemma 5.4. Clearly, $P(j, l) \leq 1$ for all j and l . By Lemma 5.4

$$P(j, l+1) \leq \frac{1}{2}P(j, l) + MP(j-1, l), \quad j \geq 1, \quad (5.24)$$

and by relation (5.15) $P(0, l) \leq e^{-\alpha 2^l D^{2/5}}$ if $l \leq L$. Hence there is a constant $k_0 > 0$ in such a way that $P(0, k_0 + l) \leq \left(\frac{2}{3}\right)^l$ if $k_0 + l \leq L$. Because of this relation, the inequality $P(j, l) \leq 1$ and formula (5.24) there is a constant $k_1 \geq k_0$ in such a way that $P(1, k_1 + l) \leq \frac{1}{3M} \left(\frac{2}{3}\right)^l$ and $P(1, k_1 + l) \leq \left(\frac{2}{3}\right)^l$ if $k_1 + l \leq L$. Similarly, there is a constant k_2 such that $P(2, k_2 + l) \leq \frac{1}{3M} \left(\frac{2}{3}\right)^l$, and $P(2, k_2 + l) \leq \left(\frac{2}{3}\right)^l$ if $k_2 + l \leq L$. This procedure can be continued, and we get a sequence $k_0 \leq k_1 \leq k_2 \leq \dots$ in such a way that the inequality $P(p, k_p + l) \leq \left(\frac{2}{3}\right)^l$ holds if $k_p + l \leq L$. The numbers k_p depend only on the parameter κ in (1.4) and the number $\bar{\eta}$ in Condition 5. The above procedure can be continued till $k_p \leq L$. In such a way we have proved that for all fixed $j \geq 0$

$$\hat{H}_{\bar{n}(T)+l}((1-\delta)^j D) \leq C(l) \left(\frac{2}{3}\right)^l,$$

if $0 \leq l \leq L$. The above relation together with formula (5.15) imply that if Condition 5 holds with a sufficiently large constant $L = L(\bar{\eta}, t)$, then an integer $k > 0$ can be chosen independently of the parameter T in such a way that

$$\hat{H}_{\bar{n}(T)+l, T}(R) \leq 2 \exp \left\{ -\frac{e^{1/\eta^3} R^2}{\bar{\eta}} \right\} \quad \text{for all } R > 0 \text{ and } k \leq l \leq L(\bar{\eta}, t). \quad (5.25)$$

Since the measure $H_{n, T}$ defined in (3.6) satisfies the relation

$$H_{\bar{n}(T)+l, T}\{x: |x| > R\} = \hat{H}_{\bar{n}(T)+l, T} \left(\sqrt{\frac{c(\bar{n}(T))}{c(\bar{n}(T)+l)}} R \right) \leq \hat{H}_{\bar{n}(T)+l, T}(\sqrt{\bar{\eta}} R)$$

relation (5.25) implies that

$$H_{\bar{n}(T)+l, T}(R) \leq 2 \exp \left\{ -e^{1/\eta^3} R^2 \right\} \quad \text{for all } R > 0, \text{ and } l^* \leq l \leq L \quad (5.26)$$

with some appropriate $l^* \geq 0$. Relation (5.26) implies in particular that $D_{\bar{n}(T)+l}^2(T) < e^{-1/\eta^2}$, i.e. $\bar{n}(T) + l$ is in the high temperature region if $l^* \leq l \leq L$.

To complete the proof of Theorem 3.1 we have to give a lower bound for $D_{\bar{n}(T)+k}^2(T)$. Let us introduce the following notation: Given two positive numbers $R_2 > R_1 > 0$ let $\mathbf{K}(R_1, R_2) = \{x: x \in \mathbb{R}^2, R_1 \leq |x| \leq R_2\}$ denote the annulus between the concentric circles with center in the origin and radii R_1 and R_2 . We claim that for any $0 \leq l \leq L$ there exist some positive numbers $R_1(l) = R_1(l, \bar{\eta}, t)$, $R_2(l) = R_2(l, \bar{\eta}, t)$ and $A(l) = A(l, \bar{\eta}, t) > 0$ such that the measure of the annulus determined by these numbers satisfies the inequality

$$\hat{H}_{\bar{n}(T)+l,T}(\mathbf{K}(R_1(l), R_2(l))) \geq A(l), \quad 0 \leq l \leq L \quad (5.27)$$

if the pair $(0, T)$ does not belong to the high temperature region. Relation (5.27) implies the required lower estimate for $D_{\bar{n}(T)+k}^2(T)$ needed in Theorem 3.1 if $k = k(T)$ is chosen as the smallest index l for which $D_{\bar{n}(T)+l}^2(T) < e^{-1/\eta^2}$. Indeed, this number k can be bounded by a number depending only on $\bar{\eta}$ and κ , and the relation between the measures $\hat{H}_{\bar{n}(T)+l,T}$ and $H_{\bar{n}(T)+l,T}$ implies that relation (5.27) also holds for $H_{\bar{n}(T)+l,T}(\mathbf{K}(R_1(k), R_2(k)))$ (i.e. the function $\hat{H}(\cdot)$ can be replaced by $H(\cdot)$ in formula (5.27)) if the radii $R_2(k)$ and $R_1(k) > 0$ are multiplied with an appropriate number. This implies that the variance $D_{\bar{n}(T)+k,T}^2$ can be bounded from below by a positive number which depends only on k and $\bar{\eta}$. We shall prove a slightly stronger statement than relation (5.27) which will be useful in later applications. We shall prove that

$$\hat{H}_{\bar{n}(T)+l,T} \left(\mathbf{K} \left(\frac{1}{2^l} R_1, \left(\frac{\sqrt{3}}{2} \right)^l R_2 \right) \right) \geq A(l), \quad 0 \leq l \leq L. \quad (5.27')$$

with some numbers $R_2 > R_1 > 0$ and $A(l) > 0$ if the pair $(0, T)$ does not belong to the high temperature region. The numbers R_j can be chosen in such a way that $R_j = R_j(\eta, \kappa)$, $j = 1, 2$.

Let us first observe that relation (5.27') holds for $l = 0$ if $\bar{n}(T)$ is not in the high temperature region. This follows from relations (5.4) and (5.5) in the case $T \leq c_0 A_0/2$ and from (5.9') and (5.11) if $T \geq c_0 A_0/2$, but $(\bar{n}(T), T)$ does not belong to the high temperature region. Indeed, formulas (5.5) and (5.11) make possible to choose the number R_2 in such a way that the $H_{\bar{n}(T),T}$ measure of the circle with center in the origin and radius $R_2 = R_2(\eta)$ is greater than $1/2$. By formulas (5.4) and (5.9') we can choose the number $R_1 = R_1(\eta)$ in such a way that by cutting out from this circle the circle with radius R_2 and center in the origin the remaining annulus $\mathbf{K}(R_1, R_2)$ has a measure greater than $1/4$. We claim that

$$\hat{H}_{\bar{n}(T)+l+1,T} \left(\mathbf{K} \left(\frac{\bar{R}_1}{2}, \frac{\sqrt{3}}{2} \bar{R}_2 \right) \right) \geq B(\bar{R}_1, \bar{R}_2, \bar{\eta}) \hat{H}_{\bar{n}(T)+l,T}(\mathbf{K}(\bar{R}_1, \bar{R}_2))^2 \quad (5.28)$$

for all $0 \leq l \leq L$ and $\bar{R}_2 > \bar{R}_1 > 0$ and an appropriate constant $B(\bar{R}_1, \bar{R}_2, \eta) > 0$. Relation (5.27') follows from (5.28) and the previous argument.

In the proof of relation (5.28) we exploit the relation

$$\begin{aligned} & \left\{ (u, x) : u \in \mathbb{R}^2, x \in \mathbb{R}^2, \frac{\bar{R}_1}{2} \leq \left| \frac{x+u}{2} \right| \leq \frac{\sqrt{3}}{2} \bar{R}_2, \frac{\pi}{3} \leq \arg(x, u) \leq \frac{\pi}{2} \right\} \\ & \supset \left\{ (u, x) : u \in \mathbb{R}^2, x \in \mathbb{R}^2, \bar{R}_1 \leq |x|, |u| \leq \bar{R}_2, \frac{\pi}{3} \leq \arg(x, u) \leq \frac{\pi}{2} \right\}. \end{aligned}$$

It follows from relation (5.14') that $Z_{\bar{n}(T)+l+1}(T) \leq 1$, since we get an upper bound for it by omitting the kernel term $\exp \left\{ -\frac{c^{(n)}}{c^{(\bar{n}(T))}} u^2 \right\}$ from the integral in (5.14'). Hence the previous relation together with (5.21) and the rotational invariance of the measure $\hat{H}_{\bar{n}(T)+l, T}$ yield that

$$\begin{aligned} \hat{H}_{\bar{n}(T)+l+1, T} \left(\mathbf{K} \left(\frac{\bar{R}_1}{2}, \frac{\sqrt{3}}{2} \bar{R}_2 \right) \right) &= \frac{1}{Z_{\bar{n}(T)+l+1}(T)} \int \int_{\frac{\sqrt{3}}{2} \bar{R}_2 \geq \left| \frac{x+u}{2} \right| \geq \frac{\bar{R}_1}{2}, x, u \in \mathbb{R}^2} \\ & \quad \exp \left\{ -\frac{c^{(\bar{n}(T)+l)}}{c^{(\bar{n}(T))}} \frac{(x-u)^2}{4} \right\} \hat{H}_{\bar{n}(T)+l, T}(dx) \hat{H}_{\bar{n}(T)+l, T}(du) \\ &\geq e^{-\bar{R}_2^2/\bar{\eta}} \int \int_{\frac{\sqrt{3}}{2} \bar{R}_2 \geq \left| \frac{x+u}{2} \right| \geq \frac{\bar{R}_1}{2}, x, u \in \mathbb{R}^2, \frac{\pi}{3} \leq \arg(x, u) \leq \frac{\pi}{2}} \hat{H}_{\bar{n}(T)+l, T}(dx) \hat{H}_{\bar{n}(T)+l, T}(du) \\ &\geq e^{-\bar{R}_2^2/\bar{\eta}} \int \int_{\bar{R}_2 \geq |x|, |u| \geq \bar{R}_1, \frac{\pi}{3} \leq \arg(x, u) \leq \frac{\pi}{2}} \hat{H}_{\bar{n}(T)+l, T}(dx) \hat{H}_{\bar{n}(T)+l, T}(du) \\ &= \frac{1}{12} e^{-\bar{R}_2^2/\bar{\eta}} \hat{H}_{\bar{n}(T)+l, T}(\mathbf{K}(\bar{R}_1, \bar{R}_2))^2. \end{aligned}$$

The last estimate implies relation (5.28) with $B(\bar{R}_1, \bar{R}_2, \bar{\eta}) = \frac{1}{12} e^{-\bar{R}_2^2/\bar{\eta}}$. Theorem 3.1 is proved.

6. Estimates in the High Temperature Region. Proof of Theorem 3.3

To study the behaviour of the function $f_n(x, T)$ in the high temperature region we need a starting index $n = \tilde{n}(T)$ for which a good estimate is known about the tail behaviour of the measure $H_{\tilde{n}(T), T}$. We also need a lower bound for the variance $D_n^2(T)$ defined in (3.5) for $n \geq \tilde{n}(T)$. This requirement will be also taken into consideration in the definition of $\tilde{n}(T)$. Let us first define the number

$$l_0 = l_0(T) = \min \left\{ l: \left(\frac{\sqrt{3}}{2} \right)^l R_2 \leq \frac{\bar{\eta}}{10} \right\} \quad (6.1)$$

if the pair $(0, T)$ is not in the high temperature region, where $\bar{\eta}$ appeared in condition 3, and the number R_2 was introduced in formula (5.27'). Now define

$$\tilde{n}(T) = \begin{cases} 0 & \text{if } (0, T) \text{ is in the high temperature region} \\ \bar{n}(T) + l \text{ with the smallest } l \text{ satisfying both (5.26) and the} \\ & \text{inequality } l \geq l_0 \text{ with } l_0 \text{ defined in (6.1)} \\ & \text{if } (0, T) \text{ is not in the high temperature region.} \end{cases} \quad (6.2)$$

It follows from the results of the previous section that for a temperature T which is not in the low temperature region the inequality $0 \leq \tilde{n}(T) - \bar{n}(T) \leq L(\bar{\eta}, t)$ holds if the number L in Condition 5 is chosen sufficiently large. The measure $H_{\tilde{n}(T), T}$ introduced in formula (3.6) is strongly concentrated around the origin. Indeed, formulas (5.12) and (5.26) give a good estimate for the $H_{\tilde{n}(T), T}$ measure of the sets $\{x: |x| \leq R\}$ for all $R \geq 0$.

Let us introduce the moments of the functions $h_{\tilde{n}(T)+l}(x, T)$ defined in (3.5).

$$M_k(l, T) = \int_{\mathbb{R}^2} |x|^k h_{\tilde{n}(T)+l}(x, T) dx \quad l \geq 0, k \geq 1.$$

We shall estimate the moments $M_2(l, T)$ and $M_4(l, T)$. It follows from relations (5.12) and (5.26) that

$$M_2(0, T) \leq \eta^* \quad \text{and} \quad M_4(0, T) \leq \eta^* \quad \text{with } \eta^* = e^{-1/\eta^2} \quad (6.3)$$

for all $T > 0$ which is not in the low temperature region. To get lower bounds for the second moments $M_2(l, T)$ let us introduce the truncated second moments

$$M_{2, \text{tr.}}(l, T) = M_{2, \text{tr.}}\left(\frac{1}{10}, l, T\right) = \int_{|x| \leq \frac{1}{10}} |x|^2 h_{\tilde{n}(T)+l}(x, T) dx. \quad (6.4)$$

It follows from (5.9'') if $(0, T)$ is in the high temperature region and from (5.27') and the definition of $\tilde{n}(T)$ if $(0, T)$ is not in the high temperature region that

$$\begin{aligned} M_{2, \text{tr.}}(0, T) &> 0, \quad \text{for all } T \geq c_0 A_0 / 2 \\ M_{2, \text{tr.}}(0, T) &> \tilde{\eta}, \quad \text{if } T \geq c_0 A_0 / 2 \text{ and } (0, T) \text{ is not in the high temperature region} \end{aligned} \quad (6.5)$$

with some $\tilde{\eta} = \tilde{\eta}(\eta, \kappa) > 0$. First we shall bound $M_2(l, T)$ and $M_4(l, T)$ from above in Lemma (6.1) for all $l \geq 0$. Then the second moment $M_2(l, T)$ will be bounded from below in Lemma (6.2). These estimates enable us to prove the central limit theorem for $g_{\tilde{n}(T)+l}(x, T)$ by means of the characteristic function technique.

Simple calculation yields that

$$M_k(l+1, T) = \frac{2}{Z_l(T)} \int e^{-u^2} |x|^k h_{\tilde{n}(T)+l} \left(\frac{x}{\sqrt{c_{\tilde{n}(T)+l+1}}} - u, T \right) h_{\tilde{n}(T)+l} \left(\frac{x}{\sqrt{c_{\tilde{n}(T)+l+1}}} + u, T \right) dx du \quad (6.6)$$

for all $l \geq 0$ and $k \geq 1$ with

$$Z_l(T) = 2 \int e^{-u^2} h_{\tilde{n}(T)+l} \left(\frac{x}{\sqrt{c_{\tilde{n}(T)+l+1}}} - u, T \right) h_{\tilde{n}(T)+l} \left(\frac{x}{\sqrt{c_{\tilde{n}(T)+l+1}}} + u, T \right) dx du. \quad (6.6')$$

These formulas will be used in the proof of the following

Lemma 6.1. *Under the conditions of Theorem 3 the inequalities*

$$M_2(l, T) \leq \eta^* \left(\frac{2}{3} \right)^l, \quad (6.7)$$

$$Z_l(T) \geq c_{\tilde{n}(T)+l} \left(1 - 6\sqrt{\eta^*} \left(\frac{5}{6} \right)^l \right), \quad (6.7')$$

$$M_2(l+1, T) \leq \frac{c_{\tilde{n}(T)+l+1}}{2} \left(1 + 10\sqrt{\eta^*} \left(\frac{5}{6} \right)^l \right) M_2(l, T), \quad (6.7'')$$

$$M_2(l, T) \leq 2 \cdot 2^{-l} \frac{c^{(\tilde{n}(T)+l)}}{c^{(\tilde{n}(T))}} \eta^* \quad \text{and} \quad M_4(l, T) \leq 5 \cdot 4^{-l} \left(\frac{c^{(\tilde{n}(T)+l)}}{c^{(\tilde{n}(T))}} \right)^2 \eta^* \quad (6.8)$$

hold for all $l \geq 0$ with the same number η^* which appears in (6.3).

Proof. Relation (6.7) holds for $l = 0$ by relation (6.3). We shall prove that if relation (6.7) holds for an integer l , then relations (6.7') and (6.7'') also hold for this l . Then we prove that if relations (6.7) and (6.7'') hold for some l , then relation (6.7) holds also for $l+1$. These statements imply relations (6.7) — (6.7''). We prove them with the help of the following calculations.

It follows from formulas (6.6) and (6.6') that

$$\begin{aligned} M_k(l+1, T) &= \frac{c_{\tilde{n}(T)+l+1}}{Z_l(T)} \int \exp \left\{ -\frac{(x-u)^2}{4} \right\} \left| \frac{x+u}{2} \right|^k c_{\tilde{n}(T)+l+1}^{k/2} \\ &\quad h_{\tilde{n}(T)+l}(x, T) h_{\tilde{n}(T)+l}(u, T) dx du \\ &\leq \frac{c_{\tilde{n}(T)+l+1}^{k/2+1}}{2^k Z_l(T)} \int |x+u|^k h_{\tilde{n}(T)+l}(x, T) h_{\tilde{n}(T)+l}(u, T) dx du \end{aligned} \quad (6.9)$$

for all $l \geq 0$ and $k \geq 1$, and

$$\begin{aligned}
Z_l(T) &= c_{\tilde{n}(T)+l+1} \int \exp \left\{ -\frac{(x-u)^2}{2} \right\} h_{\tilde{n}(T)+l}(x, T) h_{\tilde{n}(T)+l}(u, T) dx du \\
&\geq c_{\tilde{n}(T)+l+1} e^{-4\sqrt{M_2(l, t)}} \int_{|x| \leq M_2(l, t)^{1/4}, |u| \leq M_2(l, t)^{1/4}} h_{\tilde{n}(T)+l}(x, T) h_{\tilde{n}(T)+l}(u, T) dx du \\
&\geq c_{\tilde{n}(T)+l+1} e^{-4\sqrt{M_2(l, t)}} \left(1 - \frac{1}{\sqrt{M_2(l, t)}} \int_{|x| \leq M_2(l, t)^{1/4}, |u| \leq M_2(l, t)^{1/4}} x^2 h_{\tilde{n}(T)+l}(x, T) h_{\tilde{n}(T)+l}(u, T) dx du \right) \\
&\geq c_{\tilde{n}(T)+l+1} e^{-4\sqrt{M_2(l, t)}} \left(1 - \sqrt{M_2(l, t)} \right)
\end{aligned}$$

The last relation and formula (6.7) for l together imply that

$$\begin{aligned}
Z_l(T) &\geq c_{\tilde{n}(T)+l+1} \left(1 - 5\sqrt{M_2(l, t)} \right) \left(1 - \sqrt{M_2(l, t)} \right) \\
&\geq c_{\tilde{n}(T)+l+1} \left(1 - 6\sqrt{M_2(l, t)} \right) \geq c_{\tilde{n}(T)+l} \left(1 - 6\sqrt{\eta^*} \left(\frac{5}{6} \right)^l \right),
\end{aligned}$$

and this is relation (6.7') for the number l . Relation (6.9) for $k = 2$ and formula (6.7') for l together yield that

$$M_2(l+1, T) \leq \frac{c_{\tilde{n}(T)+l+1}^2}{2Z_l(T)} M_2(l, T) \leq \frac{c_{\tilde{n}(T)+l+1}}{2} \left(1 + 10\sqrt{\eta^*} \left(\frac{5}{6} \right)^l \right) M_2(l, T),$$

and this is formula (6.7'') for l . Finally, if η is chosen sufficiently small, then formulas (6.7) and (6.7'') for l imply (6.7) for $l+1$. Thus formulas (6.7) — (6.7'') are proved.

The first relation in (6.8) follows from the first relation in (6.3) and (6.7'). Formula (6.9) with the choice $k = 4$, (6.6') and the first formula in (6.8) imply that

$$\begin{aligned}
M_4(l+1, T) &\leq \frac{c_{\tilde{n}(T)+l+1}^3}{8Z_l(T)} (3M_2(l, T)^2 + M_4(l, T)) \\
&\leq \frac{1}{8} c_{\tilde{n}(T)+l+1}^2 \left(1 + 10\sqrt{\eta^*} \left(\frac{5}{6} \right)^l \right) (3M_2(l, T)^2 + M_4(l, T)) \\
&\leq 4^{-l} \left(\frac{c^{(\tilde{n}(T)+l)}}{c^{(\tilde{n}(T))}} \right)^2 \eta^{*2} + \frac{M_4(l, T)}{8}.
\end{aligned}$$

The second relation in (6.8) follows by induction from the last inequality and the second inequality in (6.3). Lemma 6.1 is proved.

Remark. The Corollary formulated after Theorem 3.1 follows from Theorem 3.1, formula (6.8) and Lemma 4.4. Indeed, if T is not in the low temperature, then by Theorem 3.1 the pair $(\tilde{n}(T), T)$ with the definition of $\tilde{n}(T)$ given in (6.1) is in the high temperature domain. By formula (6.8) all pairs (n, T) , $n \geq \tilde{n}(T)$, are in the high temperature region, i.e. if $T > 0$ is not in the low temperature region, then it is in the high temperature region. The remaining statements of the Corollary are contained in Lemma 4.4.

In the next lemma we prove an estimate from below for $M_2(l, T)$.

Lemma 6.2. *Put*

$$\sigma^2(l, T) = 2^l \frac{c^{(\tilde{n}(T))}}{c^{(\tilde{n}(T)+l)}} M_2(l, T), \quad l \geq 0. \quad (6.10)$$

Under the conditions of Theorem 3.3 the limit

$$\bar{\sigma}^2(T) = \lim_{l \rightarrow \infty} \sigma^2(l, T) > 0 \quad (6.11)$$

exists, and it is positive for all $T > 0$. If $\tilde{n}(T) \neq 0$, i.e. if $(0, T)$ is not in the high temperature region, then there exist two constants $C_2 > C_1 > 0$ depending only on the parameter $\tilde{\eta}$ in formula (6.3') in such a way that the inequalities

$$C_1 \leq \sigma^2(T) \leq C_2 \quad (6.11')$$

hold. The upper bound in (6.11') holds for all $T > 0$ which is not in the low temperature region.

Proof. The hard part of the proof is to show that $\sigma^2(l, T)$ has a non-negative $\liminf$. It follows simply from formula (6.6') that $Z_l(T) \leq c_{\tilde{n}(T)+l+1}$. A natural lower bound for $M_2(l, T)$ can be obtained in the following way. By formula (6.6) and the upper bound for $Z_l(T)$

$$\begin{aligned} M_2(l+1, T) &\geq c_{\tilde{n}(T)+l+1} \int e^{-(x-u)^2/4} \left| \frac{x+u}{2} \right|^2 h_{\tilde{n}(T)+l}(x, T) h_{\tilde{n}(T)+l}(u, T) \, dx \, du \\ &\geq \frac{c_{\tilde{n}(T)+l+1}}{4} \left(2M_2(l, T) - \int |x+u|^2 \left(1 - e^{-(x+y)^2/4} \right) \right. \\ &\quad \left. h_{\tilde{n}(T)+l}(x, T) h_{\tilde{n}(T)+l}(u, T) \, dx \, du \right) \\ &\geq \frac{c_{\tilde{n}(T)+l+1}}{2} \left(M_2(l, T) - \int \frac{1}{8} |x+u|^2 |x-u|^2 \right. \\ &\quad \left. h_{\tilde{n}(T)+l}(x, T) h_{\tilde{n}(T)+l}(u, T) \, dx \, du \right) \\ &\geq \frac{c_{\tilde{n}(T)+l+1}}{2} \left(M_2(l, T) - \int \frac{1}{2} (|x|^4 + |u|^4) \right. \end{aligned}$$

$$\begin{aligned}
& h_{\tilde{n}(T)+l}(x, T) h_{\tilde{n}(T)+l}(u, T) \, dx \, du \Big) \\
& = \frac{c_{\tilde{n}(T)+l+1}}{2} (M_2(l, T) - M_4(l, T)). \tag{6.12}
\end{aligned}$$

However, this estimate is useful only if we know that the right-hand side in it is non-negative. We do not know such an estimate for small l , hence in this case we apply a different argument. Clearly

$$M_2(l, T) \geq M_{2,\text{tr.}}(l, T),$$

where $M_{2,\text{tr.}}(l, T)$ is the truncated moment. On the other hand, we get by using an argument similar to the previous calculation and making the observation

$$\begin{aligned}
& \left\{ (x, u) : x \in \mathbb{R}^2, u \in \mathbb{R}^2, c_{\tilde{n}(T)+l+1} \left| \frac{x+u}{2} \right| \leq \frac{1}{10} \right\} \\
& \supset \left\{ (x, u) : x \in \mathbb{R}^2, u \in \mathbf{R}^2, |x| \leq \frac{1}{10}, |u| \leq \frac{1}{10}, \arg(x, u) \subset I \right\}
\end{aligned}$$

with $I = \left(\frac{\pi}{50}, \frac{49\pi}{50} \right) \cup \left(\frac{51\pi}{50}, \frac{99\pi}{50} \right)$ that

$$\begin{aligned}
M_{2,\text{tr.}}(l+1, T) & \geq c_{\tilde{n}(T)+l+1} \int_{c_{\tilde{n}(T)+l+1} \left| \frac{x+u}{2} \right| \leq \frac{1}{10}} e^{-(x-u)^2/4} \left| \frac{x+u}{2} \right|^2 \\
& \quad h_{\tilde{n}(T)+l}(x, T) h_{\tilde{n}(T)+l}(u, T) \, dx \, du \\
& \geq c_{\tilde{n}(T)+l+1} e^{-1/100} \int_{|x| \leq \frac{1}{10}, |u| \leq \frac{1}{10}, \arg(x, u) \subset I} \left| \frac{x+u}{2} \right|^2 \\
& \quad h_{\tilde{n}(T)+l}(x, T) h_{\tilde{n}(T)+l}(u, T) \, dx \, du \\
& = \frac{c_{\tilde{n}(T)+l+1}}{4} e^{-1/100} \int_{|x| \leq \frac{1}{10}, |u| \leq \frac{1}{10}, \arg(x, u) \subset I} (x^2 + u^2) \\
& \quad h_{\tilde{n}(T)+l}(x, T) h_{\tilde{n}(T)+l}(u, T) \, dx \, du \\
& = c_{\tilde{n}(T)+l+1} e^{-1/100} \frac{12}{25} M_{2,\text{tr.}}(l, T) \geq \frac{1}{3} c_{\tilde{n}(T)+l+1} M_{2,\text{tr.}}(l, T).
\end{aligned}$$

The last estimate implies that

$$\sigma^2(l, T) = 2^l \frac{c^{(\tilde{n}(T))}}{c^{(\tilde{n}(T)+l)}} M_2(l, T) \geq 2^l \frac{c^{(\tilde{n}(T))}}{c^{(\tilde{n}(T)+l)}} M_{2,\text{tr.}}(l, T) \geq \left(\frac{2}{3} \right)^l M_{2,\text{tr.}}(0, T). \tag{6.13}$$

On the other hand, it follows from (6.12) and the second inequality in (6.8) that

$$\begin{aligned}
\sigma^2(l+1, T) & \geq \sigma^2(l, T) - 2^l \frac{c^{(\tilde{n}(T))}}{c^{(\tilde{n}(T)+l+1)}} M_4(l, T) \\
& \geq \sigma^2(l, T) - \frac{5\eta^*}{2^l c_{\tilde{n}(T)+l+1}} \frac{c^{(\tilde{n}(T)+l)}}{c^{(\tilde{n}(T))}} \geq \sigma^2(l, T) - 5\eta^* \left(\frac{3}{4} \right)^l. \tag{6.14}
\end{aligned}$$

Because of (6.13) and (6.5) an index $\bar{l} \geq 0$ can be chosen in such a way that

$$\sigma^2(\bar{l}, T) \geq 100\eta^* \left(\frac{3}{4}\right)^{\bar{l}}.$$

Moreover, $\bar{l} \leq K(\bar{\eta}, \kappa)$ with some appropriate $K(\bar{\eta}, \kappa)$, if the pair $(n = 0, T)$ is not in the high temperature region. Hence, relation (6.14) implies that

$$\frac{\sigma^2(\bar{l} + l + 1, T)}{\sigma^2(\bar{l}, T)} \geq \frac{\sigma^2(\bar{l} + l, T)}{\sigma^2(\bar{l}, T)} - \frac{1}{20} \left(\frac{3}{4}\right)^l.$$

This relation together with the bound on $\sigma^2(\bar{l}, T)$ imply that $\liminf_{l \rightarrow \infty} \sigma^2(l, T) > 0$, and this liminf can be bounded by a positive number depending only on $\bar{\eta}$ and κ if $(0, T)$ is not in the high temperature region. The analogous result for limsup follows from (6.7''). To complete the proof it is enough to show that the liminf is actually lim. To prove this let us observe that for any $\varepsilon > 0$ and $N > 0$ there is some $m > N$ such that $\sigma^2(m, T) < \liminf_{n \rightarrow \infty} \sigma^2(n, T) + \varepsilon$. Then by formula (6.7'')

$$\sigma^2(n, T) \leq \sigma^2(m, T) \prod_{l=m}^n \left(1 + 10\sqrt{\eta^*} \left(\frac{5}{6}\right)^l\right) \leq \liminf_{n \rightarrow \infty} \sigma^2(n, T) + 2\varepsilon, \quad n > m$$

for any $\varepsilon > 0$ if $N = N(\varepsilon)$ is chosen sufficiently large. Lemma 6.2 is proven.

To prove Theorem 3.3 let us introduce the characteristic functions

$$\varphi_n(s, T) = \int_{\mathbb{R}^2} e^{isx} \tilde{h}_n(x, T) dx, \quad s \in \mathbb{R}^2 \quad (6.15)$$

and moments

$$\tilde{M}_k(n, T) = \int_{\mathbb{R}^2} |x|^k \tilde{h}_n(x, T) dx,$$

where the function $\tilde{h}(x, T)$ was defined in (3.24). Clearly,

$$\tilde{M}_k(n, T) = \left(\frac{2^n}{c^{(n)}}\right)^{k/2} M_k(n - \tilde{n}(T), T) \quad \text{if } n \geq \tilde{n}(T).$$

In particular, $\tilde{M}_2(n, T) = \frac{2^{\tilde{n}(T)}}{c^{(\tilde{n}(T))}} \sigma^2(n - \tilde{n}(T), T)$. We shall prove Theorem 3.3 by means of the usual characteristic function technique. The following lemma plays a crucial role in the proof.

Lemma 6.3. *Under the conditions of Theorem 3.3 the relation*

$$\lim_{n \rightarrow \infty} \frac{c^{(\tilde{n}(T))}}{2^{\tilde{n}(T)}} \tilde{M}_2(n, T) = \bar{\sigma}^2(T) \quad (6.16)$$

holds with the constant $\bar{\sigma}^2(T)$ appearing in Lemma 6.2, and

$$\lim_{n \rightarrow \infty} \sup_{|s| \leq A} \left| \log \varphi_n(s, T) + \frac{2^{\tilde{n}(T)}}{c^{(\tilde{n}(T))}} \bar{\sigma}^2(T) \frac{s^2}{2} \right| \rightarrow 0 \quad (6.17)$$

for all $A > 0$.

Proof. Relation (6.16) follows from Lemma 6.2, and it follows from the second relation in (6.8) that $\tilde{M}_4(n, T) \leq \left(\frac{2^{\tilde{n}(T)}}{c^{(\tilde{n}(T))}} \right)^2 \eta^*$. Hence the characteristic function φ can be estimated as

$$\left| \varphi_n(s, T) - \left(1 - \tilde{M}_2(n, T) \frac{s^2}{2} \right) \right| \leq \left(\frac{2^{\tilde{n}(T)}}{c^{(\tilde{n}(T))}} \right)^2 \eta^* |s|^4 \quad \text{for } n \geq \tilde{n}(T) \text{ and } s \in \mathbb{R}^2. \quad (6.18)$$

The coefficient of $|s|^4$ is bounded by a constant (depending on T), and the coefficient at $|s|^2$ converges to the positive constant $\frac{2^{\tilde{n}(T)}}{c^{(\tilde{n}(T))}} \bar{\sigma}^2(T)$. Hence formula (6.18) implies that for any $\varepsilon > 0$,

$$\left| \log \varphi_n(s, T) + \frac{2^{\tilde{n}(T)}}{c^{(\tilde{n}(T))}} \bar{\sigma}^2(T) \frac{s^2}{2} \right| \leq \varepsilon \quad \text{if } n > n_1 \text{ and } |s| \leq \delta \quad (6.19)$$

with some $n_1 = n_1(\varepsilon, T)$ and $\delta = \delta(\varepsilon, T)$. By a rescaled version of the recursive formula (2.12) we can write

$$\begin{aligned} \varphi_{n+1}(\sqrt{2}s, T) &= \frac{1}{Z_n(T)} \int \exp \left\{ is(x+u) - \frac{c^{(n)}(x-u)^2}{4 \cdot 2^n} \right\} \tilde{h}_n(x, T) \tilde{h}_n(u, T) dx du \\ &= \frac{1}{Z_n(T)} \left[\varphi_n(s, T)^2 - \int e^{is(x+u)} \left(1 - \exp \left\{ -\frac{c^{(n)}(x-u)^2}{4 \cdot 2^n} \right\} \right) \right. \\ &\quad \left. \tilde{h}_n(x, T) \tilde{h}_n(u, T) dx du \right] \end{aligned}$$

with

$$Z_n(T) = \int \exp \left\{ -\frac{c^{(n)}(x-u)^2}{4 \cdot 2^n} \right\} \tilde{h}_n(x, T) \tilde{h}_n(u, T) dx du.$$

The estimates

$$\begin{aligned} &\left| \int e^{is(x+u)} \left(1 - \exp \left\{ -\frac{c^{(n)}(x-u)^2}{4 \cdot 2^n} \right\} \right) \tilde{h}_n(x, T) \tilde{h}_n(u, T) dx du \right| \\ &\leq \int \frac{c^{(n)}(x-u)^2}{4 \cdot 2^n} \tilde{h}_n(x, T) \tilde{h}_n(u, T) dx du = \frac{c^{(n)}}{2^n} \tilde{M}_2(n, T) \end{aligned}$$

and similarly

$$1 \geq Z_n(T) \geq 1 - \frac{c^{(n)}}{2^n} \tilde{M}_2(n, T)$$

hold. Hence

$$\varphi_n^2(s, T) - \frac{c^{(n)}}{2^n} \tilde{M}_2(n, T) \leq \varphi_{n+1}(\sqrt{2}s, T) \leq \frac{\varphi_n^2(s, T) + \frac{c^{(n)}}{2^n} \tilde{M}_2(n, T)}{1 - \frac{c^{(n)}}{2^n} \tilde{M}_2(n, T)}.$$

The term $\frac{c^{(n)}}{2^n} \tilde{M}_2(n, T)$ is much less than $\left(\frac{2}{3}\right)^n$ for large n . If we have a positive lower bound on $\varphi_n(s)$ then we get, by taking logarithm in the last relation, that

$$\left| \log \varphi_{n+1}(\sqrt{2}s, T) - 2 \log \varphi_n(s, T) \right| \leq \left(\frac{2}{3}\right)^n \quad \text{if } n > n_2 \text{ and } \varphi_n(s, T) \geq \frac{1}{K} \quad (6.20)$$

with some $n_2 = n_2(K, T)$. Formula (6.17) can be deduced from (6.19) and (6.20). Indeed, define an index k by the relation $A \leq \delta 2^{k/2} < \sqrt{2}A$ with the numbers A and δ in (6.17) and (6.18). Put $K = 2e^{-2\tilde{n}(T)\bar{\sigma}^2(T)A^2/c^{(\tilde{n}(T))}}$ and let $\varepsilon \leq \frac{1}{8K}$. Choose a

number n_3 such that $\left(\frac{2}{3}\right)^{n_3} \leq \varepsilon$, and let us consider such indices n for which $n \geq \max(n_1(\varepsilon, T), n_2(K, T), n_3)$. Then simple induction yields that for $j \leq k$

$$\left| \varphi_{n+j}(s, T) + \frac{2^{\tilde{n}(T)}}{c^{(\tilde{n}(T))}} \bar{\sigma}^2(T) \frac{s^2}{2} \right| \leq \varepsilon + 3 \left(1 - \left(\frac{2}{3}\right)^{j+1} \right) \leq 4\varepsilon \quad \text{and} \quad |\varphi_{n+j}(s, T)| \geq \frac{1}{K}$$

for $j \leq k$. Since ε can be chosen arbitrary small in the last relation, it implies relation (6.17). Lemma 6.3 is proved.

Theorem 3.3 simply follows from Lemmas 6.2 and 6.3. Indeed, Lemma 6.3 implies that the measures $\tilde{H}_{n,T}$ converge in distribution to the normal law with expectation zero and covariance $\frac{2^{\tilde{n}(T)}}{c^{(\tilde{n}(T))}} \bar{\sigma}^2(T) \mathbf{I}$. The bounds obtained for the variance follow from Lemma 6.2 and the observation that the difference $\tilde{n}(T) - \bar{n}(T)$ can be bounded by a number depending only on $\bar{\eta}$ and κ .

Let us finally show that Corollary to Theorem 3.3 simply follows from Theorem 3.3. By formulas (2.5), (2.10), and (3.8) we can write

$$2^{-n} p_n(2^{-n/2} \sqrt{T} x, T) = C(n) \exp \left\{ -\frac{\bar{c}^{(n)} A_n}{2^{n+1}} x^2 \right\} \tilde{h}_n(x, T) \quad (6.21)$$

with an appropriate norming constant $C(n)$. Observe that the expressions at both sides of this identity are density functions, the measures with density function $\tilde{h}_n(x, T)$ have a limit as $n \rightarrow \infty$, the term $\left\{ -\frac{\bar{c}^{(n)} A_n}{2^{n+1}} x^2 \right\}$ is bounded, and it tends to 1 uniformly in any compact set as $n \rightarrow \infty$. These facts imply that $C(n) \rightarrow 1$ in (6.21), and the measures with density functions $2^{-n} p_n(2^{-n/2} \sqrt{T} x, T)$ have the same limit as the measures with density functions $\tilde{h}_n(x, T)$. Hence the Corollary of Theorem 3.3 holds.

7. Estimates in the Low Temperature Region. Proof of Theorem 3.2

The proof of Theorem 3.2 heavily exploits the results of Section 4. These results show that by replacing the operator $\mathbf{Q}_n$, whose application makes possible to compute the function $f_{n+1}(x, T)$, with its linearization $\mathbf{T}_n$ only a negligible error is committed. Formula (4.17) enables one to investigate the operator $\mathbf{T}_n$ in the Fourier space. In such a way good estimates can be obtained for the Fourier transform of a regularized version of the function $f_{n+1}(x, T)$. The results of Theorem 3.2 can be proved by means of these estimates with the help of inverse Fourier transformation.

Formulas (3.8) and (3.9) were already proved in Lemma 4.4. The proof of the statement that the fix point equation (1.30) has a unique non-zero solution which is a density function, is a simple adaptation of Lemma 12 in [BM3]. It is enough to observe that in that proof the parameter c which was taken there $1 < c < \sqrt{2}$ can be chosen also as $c = 1$. Also the tail behaviour of the function $g(x)$ and that of its Fourier transform $\tilde{g}(t)$ together with its analytic continuation $\tilde{g}(t + is)$ can be studied similarly to the proof of Lemma 13 in [BM3]. In such a way one gets the following inequalities:

$$\left| \frac{d^j}{dx^j} g(x) \right| \leq \begin{cases} C_j(\varepsilon) e^{-2(1-\varepsilon)|x|} & \text{for all } x, \quad j = 0, 1, 2, \dots, \\ C_j(\varepsilon, \alpha) \exp\{-Ax^\alpha\}, & \text{for } x > 0. \quad j = 0, 1, 2, \dots, \end{cases} \quad (7.1)$$

and

$$|\tilde{g}(t + is)| \leq \frac{C_j(\varepsilon)}{1 + |t|^j} \text{ for } |s| \leq 2(1 - \varepsilon) \text{ and arbitrary } t, \quad j = 1, 2, \dots \quad (7.2)$$

for all $\varepsilon > 0$ with appropriate constants $A > 0$, $C_j(\varepsilon) > 0$ and $C_j(\varepsilon, \alpha) > 0$.

It is simpler to work with an appropriately scaled version of the functions $f_n(x, T)$. Put

$$\bar{f}_n(x, T) = \frac{1}{M_n(T)} f_n\left(\frac{x}{M_n(T)}, T\right)$$

and

$$\bar{\varphi}_n(f_n(x, T)) = \frac{1}{M_n(T)} \varphi_n\left(f_n\left(\frac{x}{M_n(T)}, T\right)\right).$$

Let us also introduce the functions

$$\psi_{n+1}(f_n(x, T)) = \frac{1}{M_n(T)} \mathbf{T}_n \varphi_n\left(f_n\left(\frac{x}{M_n(T)}, T\right)\right).$$

The estimates of Proposition 4.2 and relation (4.17) can be rewritten for these new functions. We shall rewrite formulas (4.15) and (4.16) only in the case when $n > N_1(T)$ with the number $N_1(T)$ defined in formula (4.18), i.e. in the case when $\beta_n(T)$ and

$M_n^{-1}(T)$ have the same order of magnitude. In this case $M_n(T)\sqrt{\beta_{n+1}(T)} \leq 10$,

$$\begin{aligned} & \left| \frac{\partial^j}{\partial x^j} (\bar{f}_{n+1}(x, T) - \psi_{n+1}(f_n(x, T))) \right| \\ & \leq K_1 \frac{\beta_n(T)}{c^{(n)}} \left[\exp \left\{ -\frac{1}{10} \left| 2x + \frac{x^2}{c^{(n+1)}} \right| \right\} + \exp \left\{ -\frac{|x|}{5} \right\} \right] \leq K_2 \frac{\beta_n(T)}{c^{(n)}} e^{-|x|/10} \\ & \quad x > -c^{(n+1)} M_{n+1}^2(T), \quad j = 0, 1, 2 \end{aligned} \quad (7.3)$$

and

$$\left| \frac{\partial^j}{\partial x^j} \psi_{n+1}(f_n(x, T)) \right| \leq K_3 e^{-|x|/5}, \quad x \in \mathbb{R}^1, \quad j = 0, 1, 2, 3, 4 \quad (7.4)$$

with some universal constants K_1 , K_2 and K_3 . Formula (4.17) can be rewritten as

$$\tilde{\psi}_{n+1}(f_n(\xi, T)) = \tilde{\mathbf{T}}_n \tilde{\varphi}_n(f_n(M_n(T)\xi, T)) = \frac{\exp \left\{ i \frac{c_{n+1}}{4} \xi \right\}}{\sqrt{1 + i \frac{c_{n+1}}{2} \xi}} \tilde{\varphi}_n^2 \left(f_n \left(\frac{c_{n+1}}{2} \xi, T \right) \right). \quad (7.5)$$

We claim that under the conditions Theorem 3.2,

$$\lim_{n \rightarrow \infty} \sup_{x \geq -c^{(n)} M_n^2(T)} \left| \frac{\partial^j}{\partial x^j} (\bar{f}_n(x, T) - \bar{\varphi}(f_n(x, T))) \right| e^{|x|/20} = 0, \quad j = 0, 1, 2. \quad (7.6)$$

Indeed, by relations (7.3) and (7.4)

$$\left| \frac{\partial^j}{\partial x^j} \bar{f}_n(x, T) \right| \leq e^{-|x|/10}, \quad j = 0, 1, 2, \quad \text{if } x \geq -c^{(n)} M_n^2(T) \quad (7.7)$$

and $\bar{\varphi}_n(f_n(x, T))$ is the appropriate scaling of the function

$$\varphi \left(\frac{x}{\sqrt{c^{(n)}} M_n(T)} \right) f_n \left(\frac{x}{\sqrt{c^{(n)}} M_n(T)} \right).$$

Under the conditions of Theorem 3.2, formula (4.28) holds, which implies that

$$\lim_{n \rightarrow \infty} \sqrt{c^{(n)}} M_n(T) = \infty$$

This fact together with (7.7) allow us to give a good bound on the difference between the functions $\bar{\varphi}_n(f_n(x, T))$ and $\varphi \left(\frac{x}{\sqrt{c^{(n)}} M_n(T)} \right) f_n \left(\frac{x}{\sqrt{c^{(n)}} M_n(T)} \right)$. Relation (7.6) can be deduced from this bound and formula (7.7).

It follows from Lemma 4.4 that $\lim_{n \rightarrow \infty} \frac{M_{n+1}(T)}{M_n(T)} = 1$. Relations (7.3) and (7.6) together with this fact imply that

$$\lim_{n \rightarrow \infty} \sup_{|x| < \infty} \left| \frac{\partial^j}{\partial x^j} (\psi_n(f_{n-1}(x, T)) - \bar{\varphi}_n(f_n(x, T))) \right| e^{|x|/20} = 0, \quad j = 0, 1, 2. \quad (7.8)$$

The fix point equation (3.5) can be rewritten for the Fourier transform of the function $g_1(x) = g\left(x - \frac{1}{4}\right)$ as

$$\tilde{g}_1(\xi) = \frac{\exp\left\{\frac{i}{4}\xi\right\}}{\sqrt{1 + \frac{i}{2}\xi}} \tilde{g}_1^2\left(\frac{\xi}{2}\right). \quad (7.9)$$

(We work with the function $g_1(x)$ instead of $g(x)$ because $\int g_1(x) dx = 0$.) The right-hand side of formulas (7.5) and (7.9) are very similar. Let us recall that by Condition 1 $c_n \rightarrow 1$ as $n \rightarrow \infty$.

Now we prove, using an adaptation of the proof of Lemmas 14 and 15 in [BM3], that the Fourier transforms of the functions $\psi_{n+1}(f_n(x, T))$ converge to the Fourier transform of the function $g_1(x)$, and this convergence is uniform in all compact domains. First we prove a modified version of this statement, where ψ_n is replaced with $\bar{\varphi}_n$ in a small neighbourhood of the origin. We want to work with the functions $\log \tilde{\varphi}_n(f_n(\xi, T))$. To do this, observe first that for $n > N_1(T)$ there is some constant $A > 0$ such that all functions $\tilde{\varphi}_{n+1}(f_n(\xi, T))$ are separated from zero in the interval $|\xi| \leq A$. Indeed,

$$\begin{aligned} |1 - \tilde{\varphi}_n(f_n(\xi, T))| &\leq \int |e^{ix\xi} - 1| \bar{\varphi}_n(f_n(x, T)) dx \\ &\leq \int |\xi| |x| \bar{\varphi}_n(f_n(x, T)) dx \leq \text{const. } |\xi|. \end{aligned}$$

Similarly,

$$\left| \frac{\partial^j}{\partial \xi^j} \tilde{\varphi}_n(f_n(\xi, T)) \right| \leq C(j) \quad \text{for all } j \geq 0 \text{ and } n \geq N_1(T).$$

Hence a constant $A > 0$ can be chosen in such a way that

$$\sup_{|\xi| \leq 2A} \max \left(|1 - \tilde{g}_1(\xi)|, \sup_{n \geq N_1(T)} |1 - \tilde{\varphi}_n(f_n(\xi, T))| \right) \leq \frac{1}{2}.$$

These estimates imply that

$$\sup_{|\xi| \leq A} \sup \left| \frac{\partial^2}{\partial \xi^2} \log \tilde{\varphi}_n(f_n(\xi, T)) \right| \leq C(T) \quad (7.10)$$

with a constant $C(T) < \infty$ independent of n . We claim that

$$\sup_{|\xi| \leq A} \left| \frac{\partial^2}{\partial \xi^2} \log \tilde{\varphi}_n(f_n(\xi, T)) - \frac{d^2}{d^2 \xi} \log \tilde{g}_1(\xi) \right| \rightarrow 0 \quad \text{as } n \rightarrow \infty. \quad (7.11)$$

To prove (7.11) let us first observe that $\lim_{n \rightarrow \infty} c_n = 1$ by Condition 1. By (7.8),

$$\lim_{n \rightarrow \infty} \sup_{|\xi| \leq A} \left| \frac{\partial^2}{\partial \xi^2} \left(\log \tilde{\psi}_{n+1}(f_n(\xi, T)) - \log \tilde{\varphi}_{n+1}(f_{n+1}(\xi, T)) \right) \right| = 0,$$

and because of the estimates obtained for the derivatives of $\tilde{\varphi}_n(\xi, T)$

$$\left| \frac{\partial^2}{\partial \xi^2} \log \tilde{\varphi}_n(f_n(\xi_1, T)) - \frac{\partial^2}{\partial \xi^2} \log \tilde{\varphi}_n(f_n(\xi_2, T)) \right| \leq \text{const.} \cdot |\xi_1 - \xi_2|$$

if $|\xi_1| \leq A$ and $|\xi_2| \leq A$

for all large indices n with a constant independent of n . Taking logarithm and then differentiating twice in formulas (7.5) and (7.9) we get with the help of the above observations that

$$\begin{aligned} & \sup_{|\xi| \leq A} \left| \frac{\partial^2}{\partial \xi^2} \log \tilde{\varphi}_{n+1}(f_{n+1}(\xi, T)) - \frac{d^2}{d^2 \xi} \log \tilde{g}_1(\xi) \right| \\ & \leq \frac{1}{2} \sup_{|\xi| \leq A} \left| \frac{\partial^2}{\partial \xi^2} \log \tilde{\varphi}_n(f_n(\xi, T)) - \frac{d^2}{d^2 \xi} \log \tilde{g}_1(\xi) \right| + \delta_n(T) \end{aligned}$$

with some sequence $\lim_{n \rightarrow \infty} \delta_n(T) = 0$. This relation together with (7.10) imply (7.11). Since

$$\left. \frac{\partial}{\partial \xi} \log \tilde{\varphi}_n(f_n(\xi, T)) \right|_{\xi=0} = \left. \frac{d}{d \xi} \log \tilde{g}_1(\xi) \right|_{\xi=0} = 0 \text{ and } \log \tilde{\varphi}_n(f_n(0, T)) = \log \tilde{g}_1(0) = 0,$$

relation (7.11) also implies that

$$\lim_{n \rightarrow \infty} \sup_{|\xi| \leq A} |\tilde{\varphi}_n(f_n(\xi, T)) - \tilde{g}_1(\xi)| = 0. \quad (7.12)$$

Moreover, relation (7.12) holds for all $A > 0$. This can be proved similarly to the argument of Lemma 15 in [BM3]. One has to observe that because of the structure of formulas (7.5) and (7.9), the relation $c_n \rightarrow 1$ as $n \rightarrow \infty$, the continuity of the function $\tilde{g}_1(\xi)$ and the relation

$$\lim_{n \rightarrow \infty} \sup_{|\xi| < \infty} |\tilde{\psi}_{n+1}(f_n(\xi, T)) - \tilde{\varphi}_{n+1}(f_{n+1}(\xi, T))| = 0,$$

which also follows from (7.9), the validity of relation (7.12) in an interval $|\xi| \leq A$ also implies its validity in the interval $|\xi| \leq (2 - \varepsilon)A$ for any $\varepsilon > 0$. In relation (7.12) the function $\tilde{\varphi}_n(f_n(\xi, T))$ can be replaced by $\tilde{\psi}_{n+1}(f_n(\xi, T))$, i.e. the relation

$$\lim_{n \rightarrow \infty} \sup_{|\xi| \leq A} |\tilde{\psi}_{n+1}(f_n(\xi, T)) - \tilde{g}_1(\xi)| = 0 \quad (7.12')$$

holds for all $A > 0$. It can be proved from (7.12') by means of inverse Fourier transformation that

$$\lim_{n \rightarrow \infty} \sup_{|x| < \infty} \left| \frac{\partial^j}{\partial x^j} \psi_{n+1}(f_n(x, T)) - \frac{d^j}{dx^j} g_1(x) \right| = 0 \quad j = 0, 1, 2. \quad (7.13)$$

To prove 7.13 we need, beside the estimate (7.12'), some bound about the decrease of the functions $\tilde{g}_1(\xi)$ and $\tilde{\psi}_{n+1}(f_n(\xi, T))$ as $\xi \rightarrow \pm\infty$. The estimate (7.2) gives a good bound for the Fourier transform of the function $g_1(x)$. We can get a good estimate for the Fourier transform of $\psi_{n+1}(f_n(x, T))$ with the help of the inductive hypothesis $J(n)$ in Section 4 and relation (7.5). Rewriting the inductive hypothesis $J(n)$ for the function $\bar{\varphi}_n(f_n(x, T))$ we get with the help of some standard calculation that the Fourier transform $\tilde{\psi}_{n+1}(f_n(\xi, T))$ decreases at infinity faster than $|\xi|^{-4}$. These estimates are sufficient for the proof of (7.13). Relation (7.13) and (7.3) together give an estimate on the function $\tilde{f}_n(x, T)$ and its derivatives which is equivalent to (3.10). Theorem 3.2 is proved.

8. Estimates Near the Critical Point. Proof of Theorems 3.4, 1.3, and 1.5

Our previous results suggest that $M_{n+1}^2(T) \sim M_n^2(T) - \frac{1}{2c^{(n)}}$, hence the derivative $\frac{dM_n^2(T)}{dT}$, as a function of n , changes very little if the pair (n, T) is in the low domain region (observe that $c^{(n)}$ does not depend on T). Therefore, it is natural to expect that $\frac{M_\infty^2(T)}{dT}$ is of constant order below the critical value $T_{cr.}$, and $M_\infty^2(T) - M_\infty^2(T_{cr.}) \sim \text{const.} (T_{cr.} - T)$ for $T < T_{cr.}$. If T_n denotes the smallest T for which the pair (n, T) leaves the low temperature region at the n -th step, then the following heuristic argument may suggest the magnitude of $T_n - T_{n+1}$ for large n . Since both $c^{(n)} M_n^2(T_n) \sim \eta^{-1}$ and $c^{(n+1)} M_{n+1}^2(T_{n+1}) \sim \eta^{-1}$, beside this $M_{n+1}^2(T_{n+1}) - M_n^2(T_{n+1}) \sim \frac{1}{2c^{(n)}}$, $M_n^2(T_{n+1}) - M_n^2(T_n) \sim \frac{1}{c^{(n)}}$. On the other hand, $M_n^2(T_{n+1}) - M_n^2(T_n) \sim T_{n+1} - T_n$. This argument suggests that $T_{n+1} - T_n \sim \frac{1}{c^{(n)}}$ and $T_n - T_{cr.} \sim \sum_{k=n}^{\infty} \frac{1}{c^{(k)}}$. In this section we prove the results obtained by means of the above heuristic argument hold if the sequence $c^{(n)}$ satisfies some regularity conditions. The proofs are based on the following Theorem A.

Theorem A. *There exists $\kappa_0 = \kappa_0(N)$ such that if (i) $0 < \kappa < \kappa_0$ in formula (1.17), (ii) Conditions 1 — 4 are satisfied, (iii) $0 < \bar{T} < c_0 A_0 / 2$, and (iv) integer $n \geq 1$ such that the pair $(n, \bar{T})$ belongs to the low temperature region, then for all $0 < T < \bar{T}$ the pair (n, T) also belongs to the low temperature region, and the following inequalities hold for $T \leq \bar{T}$:*

a.) If $0 \leq n \leq N$, then

$$\frac{C_1}{\sqrt{\kappa} T^2} < -\frac{dM_{n+1}(T)}{dT} < \frac{C_2}{\sqrt{\kappa} T^2} \quad \text{with some } \infty > C_2 > C_1 > 0.$$

b.) If $n \geq N$, then

$$\frac{dM_{n+1}(T)}{dT} = \frac{dM_n(T)}{dT} \left(1 + \frac{1}{4c^{(n)}M_n^2(T)} + \frac{\delta_n(T)}{c^{(n)}} \right),$$

where $|\delta_n(T)| \leq C \frac{\beta_{n+1}(T)}{c^{(n+1)}} \beta_n(T)$ with some appropriate $C > 0$.

We shall prove Theorem A in Appendix A below. This result can be interpreted in an informal way as the “differentiation” of the asymptotic identity (4.13). In this formal differentiation we bound the absolute value of $\frac{d\sqrt{\beta_n(T)}}{dT}$ by $\text{const.} \left| \frac{M_n(T)}{dT} \right| \beta_n(T)$. The main difficulty in the proof of Theorem A is to bound the error caused by the linear approximation of the operator $\mathbf{Q}_n$ by $\mathbf{T}_n$ when differentiating with respect to T . To overcome this difficulty we need a good control not only on the functions $f_n(x, T)$ but also on their derivatives $\frac{\partial}{\partial T} f_n(x, T)$. Hence we have to work out the estimation of these derivatives. In particular, we have to find the inductive hypotheses describing their behaviour. These are the analogs of the inductive hypotheses $I(n)$ and $J(n)$ formulated in Section 4. It demands fairly much work to work out the details, but after the formulation and proof of these inductive hypotheses the proof of Theorem A is simple.

Proof of Theorem 3.4. We prove with the help of Theorem A that if the conditions of Theorem 3.4 hold, $0 < \bar{T} < c_0 A_0/2$ and the pair $(n-1, \bar{T})$ belongs to the low temperature, then there exist some constants $0 < C_1 < C_2$ independent of T such that

$$\frac{C_1}{\kappa T^3} < -\frac{dM_n^2(T)}{dT} < \frac{C_2}{\kappa T^3}. \quad (8.1)$$

for all $0 < T \leq \bar{T}$.

For $0 \leq n \leq N$ formula (8.1) follows Part a) of Theorem A and relations (4.1), (4.4) and (4.6) which give a bound on $M_n(T)$ in the case $0 \leq n \leq N$. To prove formula (8.1) for $n > N$ first we show that

$$\begin{aligned} -\frac{dM_n^2(T)}{dT} \exp \left\{ -K \left(\frac{\beta_{n+1}(T)}{c^{(n)}} \right)^2 \right\} &\leq -\frac{dM_{n+1}^2(T)}{dT} \\ &\leq -\frac{dM_n^2(T)}{dT} \exp \left\{ K \left(\frac{\beta_{n+1}(T)}{c^{(n)}} \right)^2 \right\} \end{aligned} \quad (8.2)$$

for all $T < \bar{T}$ and $n \geq N$ with an appropriate $K > 0$. Relation (8.2) is a consequence of Part b.) of Theorem A, formula (4.13), the inequality $\beta_{n+1} M_n^2(T) \geq 10$ and the relation $\frac{\beta_{n+1}(T)}{c^{(n)}} \leq \eta$ if (n, T) is in the low temperature domain. Indeed, these relations imply

that

$$\begin{aligned}
& - \left(1 - \frac{M_n(T)}{4c^{(n)}} - C_1 \frac{\beta_{n+1}^{3/2}(T)}{c^{(n)^2} M_n(T)} \right) \left(1 + \frac{1}{4c^{(n)} M_n^2(T)} - C \frac{\beta_{n+1}^2(T)}{(c^{(n)})^2} \right) \frac{dM_n^2(T)}{dT} \\
& \leq - \frac{dM_{n+1}^2(T)}{dT} \\
& \leq - \left(1 - \frac{M_n(T)}{4c^{(n)}} + C_1 \frac{\beta_{n+1}^{3/2}(T)}{c^{(n)^2} M_n(T)} \right) \left(1 + \frac{1}{4c^{(n)} M_n^2(T)} + C \frac{\beta_{n+1}^2(T)}{c^{(n)^2} M_n^2(T)} \right) \frac{dM_n^2(T)}{dT}.
\end{aligned}$$

The left and right-hand side of this inequality can be bounded by

$$- \left(1 \pm K \frac{\beta_{n+1}^2(T)}{c^{(n)^2} M_n^2(T)} \right) \frac{dM_n^2(T)}{dT},$$

and formula (8.2) can be deduced from these relations. For $N \leq n \leq N_1(T)$ with the number $N_1(T)$ defined in relation (4.18) relation (8.1) follows from (8.2) and (4.19). Since by (4.20) $\beta_{n+1} M_n^2(T) \leq 100$ if $n \geq N_1(T)$ and the pair (n, T) is in the low temperature domain, to prove formula (8.1) with the help of (8.2) for $n > N_1(T)$ it is enough to show that

$$\sum_{k=N_1(T)}^n \frac{1}{(c^{(k)} M_k^2(T))^2} \leq L \text{ if } n \geq N_1(T) \text{ and } (n, T) \text{ is in the low temperature domain}$$

with a constant $L > 0$ independent of T and n . Since $M_n^2(T) \geq \frac{1}{10\beta_{n+1}(T)} \geq \frac{1}{10\eta c^{(n)}}$

$$\text{and } M_k^2(T) = M_n^2(T) + (M_k^2(T) - M_n^2(T)) \geq \frac{1}{10\eta c^{(n)}} + \sum_{j=k}^{n-1} \frac{1}{8c^{(j)}},$$

$$\sum_{k=N_1(T)}^n \frac{1}{(c^{(k)} M_k^2(T))^2} \leq \text{const.} \sum_{k=N_1(T)}^n \frac{1}{\left(c^{(k)} \sum_{j=k}^n \frac{1}{c^{(j)}} \right)^2} \leq L$$

because of Condition 3. Hence formula (8.1) holds.

It follows from (1.3), Condition 4, and the results of Section 4 that all $T > c_0 A_0/4$ belong to the high temperature region. Indeed, it follows from formulas (4.26), (4.26'), (4.1), (4.4) and (4.10), that if $T > 0$ is in the low temperature region, then

$$0 \leq M_n^2(T) \leq M_N^2(T) - 30(M_N(T) + 1) - \sum_{n=1}^{\infty} \frac{1}{8c^{(n)}} \leq \frac{3}{\kappa T^2} - \sum_{n=1}^{\infty} \frac{1}{8c^{(n)}}$$

for all $n \geq N$, and $T \leq \left(\sum_{n=1}^{\infty} \frac{\kappa}{24c^{(n)}} \right)^{-1/2}$. Hence Condition 4 implies that $T \leq c_0 A_0/4$.

It follows from (8.2) that for fixed n the functions $M_n^2(T)$ is a strictly monotone decreasing, hence a simple induction with respect to n yields that the function $\beta_n(T)$ is a monotone increasing, continuous function of T . Put

$$T_n = \sup\{T: (T, n) \text{ is in the low temperature region}\}.$$

The sequence T_n is monotone decreasing, hence the limit $T_{\text{cr.}} = \lim_{n \rightarrow \infty} T_n$ exists, and by Lemma 4.2 $T_{\text{cr.}} > 0$ under Dyson's condition (1.3). We want to show that

$$C_1 \sum_{k=n}^{\infty} \frac{1}{c^{(k)}} \leq T_n - T_{\text{cr.}} \leq C_2 \sum_{k=n}^{\infty} \frac{1}{c^{(k)}}. \quad (8.3)$$

Since we can handle the sequence $M_n(T)$ better than the sequence $\beta_n(T)$ we also introduce the sequence $T(n)$

$$T(n) = \sup \left\{ T: M_n^2(T) \geq \frac{100}{c^{(n)}\eta} \right\}.$$

We will show that

$$T_{n+K} \leq T(n) \leq T_n \quad (8.4)$$

for all sufficiently large n with an appropriate $K > 0$, and

$$\frac{C_1}{c^{(n)}} \leq T(n) - T(n+1) \leq \frac{C_2}{c^{(n)}} \quad (8.5)$$

with some appropriate $C_2 > C_1 > 0$ for all sufficiently large n . Because of Condition 5 relation (8.3) follows from (8.4) and (8.5) together with the relation $\lim_{n \rightarrow \infty} T_n = T_{\text{cr.}}$.

If $T \leq T(n)$, and $m \leq n$ then either $m \leq N_1(T)$ with the number $N_1(T)$ defined in (4.19) or $\beta_{m+1}(T) \leq \frac{100}{M_m^2(T)} \leq \frac{100}{M_n^2(T)} \leq c^{(n)}\eta$. This implies that for $T \leq T(n)$ the pair (m, T) is in the low temperature region for all $m \leq n$, and $T(n) \leq T_n$. This is the right-hand side of relation (8.4).

To prove its left-hand side observe that because of Condition 5 there is some K such that

$$\sum_{k=n}^{n+K-1} \frac{1}{8c^{(k)}} > \frac{100}{c^{(n)}\eta}$$

for all sufficiently large n with appropriate $K > 0$. We claim that $T \geq T(n)$ the pair $(n+K, T)$ is not in the low temperature region. This relation implies the left-hand side of (8.4). If $(n+K, T)$ were in the low temperature region, then we would get with the help of formula (4.26) that

$$M_{n+K}^2(T) \leq M_n^2(T) - \sum_{k=n}^{n+K-1} \frac{1}{8c^{(k)}} < \frac{100}{c^{(n)}\eta} - \sum_{k=n}^{n+K-1} \frac{1}{8c^{(k)}} < \frac{100}{c^{(n)}\eta} - \frac{100}{c^{(n)}\eta} = 0,$$

and this is a contradiction.

To prove formula (8.5) let us first observe that because of the continuity and strict monotonicity of the function $M_n^2(T)$, $M_n^2(T(n)) = \frac{100}{c^{(n)}\eta}$. It follows from the last statement of Lemma 4.3 and formula (8.1) that $N_1(T) \leq n$ for all $T(n) - \varepsilon < T < T(n)$ with an appropriately small $\varepsilon > 0$. (The number $N_1(T)$ was defined in (4.18).). Hence we get with the help of formula (8.1) that for sufficiently large n and $T(n) - \varepsilon < T < T(n)$

$$\begin{aligned} \frac{100}{c^{(n)}\eta} - \frac{2}{c^{(n)}} + \bar{C}_1(T(n) - T) &\leq \frac{100}{c^{(n+1)}\eta} - \frac{1}{c^{(n)}} + \bar{C}_1(T(n) - T) \\ &\leq M_{n+1}^2(T) \\ &\leq \frac{100}{c^{(n)}\eta} - \frac{1}{8c^{(n)}} + \bar{C}_2(T(n) - T) \leq \frac{100}{\eta c^{(n+1)}} - \frac{1}{9c^{(n)}} + \bar{C}_2(T(n) - T) \end{aligned}$$

with some appropriate constants $\bar{C}_2 > \bar{C}_1 > 0$. Hence the solution of the equation $M_{n+1}^2(T) = \frac{100}{c^{(n+1)}\eta}$ satisfies the inequality $K_1 < c^{(n)}(T - T(n)) < K_2$ with appropriate constants $K_2 > K_1 > 0$. Since the solution of this equation is $T(n+1)$, this fact implies relation (8.5).

It is not difficult to see that $T_{\text{cr.}}$ is in the low temperature region. Since the inequality $M_n^2(T_{\text{cr.}}) \leq \text{const.} (T(n) - T_{\text{cr.}}) + \frac{100}{c^{(n)}\eta}$ holds for all large n , $\lim_{n \rightarrow \infty} M_n(T_{\text{cr.}}) = 0$. Then relation (8.1) implies that

$$C_1(T_{\text{cr.}} - T) \leq M_n^2(T_{\text{cr.}}) - M_n^2(T) \leq C_2(T_{\text{cr.}} - T)$$

with some positive constants $C_2 > C_1 > 0$ if $T_{\text{cr.}} \geq T \geq T_{\text{cr.}} - \varepsilon$. Letting n tend to infinity in the last relation we get formula (3.28). Since formula (8.3) is equivalent to (3.27) Theorem 3.4 is proved.

Proof of Theorem 1.3. By Corollary of Theorem 3.1, if the Dyson condition (1.3) is violated then all temperatures $T > 0$ belong to the high temperature region. By Corollary of Theorem 3.3, (1.24) holds and the measures $\bar{\nu}_{n,T}(dx)$ approaches the standard normal distribution as $n \rightarrow \infty$. Theorem 1.3 is proved.

Proof of Theorem 1.5, Part 1). The convergence of $\bar{\nu}_{n,T}(dx)$ to a standard Gaussian distribution and relation (1.32) follow from Corollary of Theorem 3.3. The asymptotics (1.33) follows from (3.26) and (3.27).

Part 2). Formula (1.34) follows from (3.9), and the convergence of $\bar{\nu}_{n,T_c}(dx)$ to the uniform distribution on the sphere follows from Theorems 3.2 and 3.4. Namely, Theorem 3.4 tells us that the critical temperature T_c belongs to the *low temperature region*. Then formula (3.10) proves that the probability distribution $\bar{\nu}_{n,T_c}(dx)$ converges to the uniform distribution on the sphere. As a matter of fact, (3.10) proves much more: it proves the convergence at $T = T_c$ of the distribution of normalized fluctuations of the average spin along the radius to a limit.

Part 3). By (2.10), (2.14), and (2.17),

$$p_n(x, T) = L_n^{-1}(T) \exp\left(-\frac{A_n l_n |x|^2}{2T}\right) f_n\left(\frac{c^{(n)}}{\sqrt{T}}\left(|x| - \sqrt{T} M_n(T)\right), T\right) \quad (8.6)$$

Let us write that $|x|^2 = \left(\sqrt{T} M_n(T) + |x| - \sqrt{T} M_n(T)\right)^2$, hence

$$\begin{aligned} \exp\left(-\frac{A_n l_n |x|^2}{2T}\right) &= \exp\left\{-\frac{A_n l_n}{2T} [T M_n^2(T) + 2\sqrt{T} M_n(T)(|x| - \sqrt{T} M_n(T)) \right. \\ &\quad \left. + (|x| - \sqrt{T} M_n(T))^2]\right\}, \end{aligned} \quad (8.7)$$

and substitute it into (8.6). This leads to the equation

$$p_n(x, T) = \tilde{L}_n^{-1}(T) \tilde{f}_n\left(\frac{|x| - \tilde{M}_n(T)}{\tilde{V}_n(T)}, T\right), \quad (8.8)$$

where

$$\begin{aligned} \tilde{M}_n(T) &= \sqrt{T} M_n(T), \quad \tilde{V}_n(T) = \frac{\sqrt{T}}{c^{(n)} M_n(T)} \\ \tilde{f}_n(t, T) &= f_n\left(\frac{t}{M_n(T)}, T\right) \exp\left(-\frac{A_n l_n t}{c^{(n)}} - \varepsilon_n(t, T)\right) \\ \varepsilon_n(t, T) &= \frac{A_n l_n t^2}{2(c^{(n)})^2 M_n^2(T)}. \end{aligned} \quad (8.9)$$

Observe that by (2.30) and (2.31),

$$\lim_{n \rightarrow \infty} \frac{A_n l_n}{c^{(n)}} = \frac{2}{3}, \quad \lim_{n \rightarrow \infty} \frac{A_n l_n}{2(c^{(n)})^2 M_n^2(T)} = 0, \quad (8.10)$$

hence (3.10) implies that there is some $C_0 > 0$ such that

$$\lim_{n \rightarrow \infty} \left\| \tilde{f}_n(t, T) - C_0 g\left(t - \frac{r-1}{4}\right) e^{-2t/3} \right\|' = 0, \quad (8.11)$$

where

$$\|f(t)\|' = \sum_{j=0}^2 \sup_{t \geq -c^{(n)} M_n^2(T)} e^{|t|/3} \left| \frac{d^j f(t)}{dt^j} \right|. \quad (8.12)$$

It remains to shift $\tilde{f}_n(t)$ to secure the mean value to be zero. Define

$$\tilde{\pi}_n(t, T) = C_n(T) \tilde{f}_n(t - a_0, T), \quad a_0 = a - \frac{r-1}{4}, \quad (8.13)$$

where $C_n(T)$ is a norming factor such that

$$\int_{\mathbb{R}^1} \tilde{\pi}_n(t) dt = 1. \quad (8.14)$$

Then

$$p_n(x, T) = L_n^{-1}(T) \tilde{\pi}_n \left(\frac{|x| - M_{n0}(T)}{\tilde{V}_n(T)}, T \right) \quad \text{with } M_{n0}(T) = \tilde{M}_n + a_0 \tilde{V}_n(T) \quad (8.15)$$

and

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}^1} t \tilde{\pi}_n(t) dt = 0. \quad (8.16)$$

Comparing these formulae with (1.29) we obtain (1.36), (1.38), and (1.39) with

$$M(T) = \sqrt{T} M_\infty(T), \quad \gamma(T) = \frac{\sqrt{T}}{3M_\infty(T)} = \frac{T}{3M(T)}, \quad (8.17)$$

where $M_\infty(T)$ is the limit (3.4). The estimates (8.17) follow from (3.28). Theorem 1.5 is proved.

Appendix A. Proof of Theorem A

To prove Theorem A we need good estimates on the partial derivatives $g_n(x, T) = \frac{\partial}{\partial T} f_n(x, T)$, on the derivatives of a scaled version of the functions $q_n(x, T)$. This can be done similarly to the estimation of the functions $f_n(x, T)$, done in Section 4. First we give estimates for the starting function $g_0(x, T)$, then prove that similar estimates hold for small indices n , more explicitly for $n \leq N$ with the index N defined in (1.22). Then inductive hypotheses can be formulated and proved for the functions $g_n(x, T)$. In Section 4 we have introduced certain operators $\bar{\mathbf{Q}}_n$, their normalization $\mathbf{Q}_n$ and the linearization of these operators denoted by $\bar{\mathbf{T}}_n$ and $\mathbf{T}_n$. The inductive hypotheses formulated there were closely related to the properties of these operators. Now we want to work similarly. To do this we have to introduce some new operators. We introduce certain operators $\bar{\mathbf{R}}_n$ and $\mathbf{R}_n$ which are the derivatives of the operators $\bar{\mathbf{Q}}_n$ and $\mathbf{Q}_n$ with respect to the variable T . We also need their linear approximation which we shall denote by $\bar{\mathbf{U}}_n$ and $\mathbf{U}_n$. We have to study the action of these operators on the functions $g_n(x, T) = \frac{\partial}{\partial T} f_n(x, T)$ and their Fourier transform. Although the proofs are not hard, it demands much time to work out the details even if only a brief explanation is given as in this Appendix.

An appropriate description of the asymptotic behaviour of the starting functions $f_0(x, T)$ and numbers $M_0(T)$ were already given in formulas (4.2) — (4.7). Some more calculation yields, with the help of some formulas in Section 4, the following estimates

for the derivatives of the magnetization $M_0(T)$ and the norming constant $Z_0(T)$ if $T < c_0 A_0/2$.

$$\left| \frac{d}{dT} \left(M_0(T) - \hat{M}_0(T) \right) \right| \leq \text{const.} \sqrt{\kappa}.$$

$$\frac{C_1}{\sqrt{\kappa} T^2} < -\frac{dM_0(T)}{dT} < \frac{C_2}{\sqrt{\kappa} T^2} \quad \text{with some } \infty > C_2 > C_1 > 0.$$

and

$$\left| \frac{dZ_0(T)}{dT} - \frac{\sqrt{\pi}}{2(A_0 - T)^{3/2}} \right| \leq \text{const.} \sqrt{\kappa}.$$

The derivatives of the functions $\bar{q}_0(x, T)$ and $f_0(x, T)$ satisfy the inequalities

$$\left| \frac{\partial q_0(x + M_0(T), T)}{\partial T} - \frac{\sqrt{A_0 - T}}{\sqrt{\pi}} \left(x^2 - \frac{1}{2(A_0 - T)} \right) e^{-(A_0 - T)x^2} \right| \leq \text{const.} \kappa^{1/4}$$

if $|x| < \log \kappa^{-1}$,
(A1)

and

$$\left| \frac{\partial q_0(x + M_0(T), T)}{\partial T} \right| \leq C \exp \left\{ -\frac{(A_0 - T)}{4} \left| 2x + \frac{x^2}{M_0^2(T)} \right| \right\} \quad \text{for } x \geq -M_0(T). \quad (\text{A2})$$

We shall apply the notation

$$g_n(x, T) = \frac{\partial f_n(x, T)}{\partial T}, \quad n = 0, 1, \dots \quad (\text{A3})$$

Since $f_0(x, T) = q_0(x + M_0(T), T)$ the previous estimates together with the results of Section 4 yield a sufficiently good control on $g_0(x, T)$. The functions $g_n(x, T)$, $n = 1, 2, \dots$, can be estimated inductively with respect to the parameter n .

Put

$$\bar{\mathbf{R}}_n f_n(x, T) = \frac{\partial}{\partial T} \bar{\mathbf{Q}}_{n, M_n(T)}^c f_n(x, T)$$

and

$$\mathbf{R}_n f_n(x, T) = \frac{\partial}{\partial T} \mathbf{Q}_n f_n(x, T) = g_{n+1}(x, T).$$

Then

$$\bar{\mathbf{R}}_n f_n(x, T) = \bar{\mathbf{R}}_n^{(1)} f_n(x, T) + \bar{\mathbf{R}}_n^{(2)} f_n(x, T)$$

with

$$\bar{\mathbf{R}}_n^{(1)} f_n(x, T) = 2 \int \exp \left\{ -\frac{u^2}{c(n)} - v^2 \right\} f_n(\ell_{n, M_n(T)}^+(x, u, v), T) \\ g_n(\ell_{n, M_n(T)}^-(x, u, v), T) du dv,$$

where the functions $g_n(x, T)$ and $\ell_{n, M_n(T)}^\pm(x, u, v), T)$ were defined in (A3) and (2.9), and

$$\bar{\mathbf{R}}_n^{(2)} f_n(x, T) = -2 \int \exp \left\{ -\frac{u^2}{c^{(n)}} - v^2 \right\} f_n(\ell_{n, M_n(T)}^+(x, u, v), T) h_n(x, u, v, T) \frac{\partial}{\partial x} f_n(\ell_{n, M_n(T)}^-(x, u, v), T) du dv$$

with

$$\begin{aligned} h_n(x, u, v, T) &= -\frac{\partial \ell_{n, M_n(T)}^-(x, u, v)}{\partial T} \\ &= \frac{M'_n(T) v^2}{\sqrt{\left(M_n(T) + \frac{x}{c^{(n+1)}} - \frac{u}{c^{(n)}}\right)^2 + \frac{v^2}{c^{(n)}}}} \\ &\quad \frac{1}{\sqrt{\left(M_n(T) + \frac{x}{c^{(n+1)}} - \frac{u}{c^{(n)}}\right)^2 + \frac{v^2}{c^{(n)}}} + \left(M_n(T) + \frac{x}{c^{(n+1)}} - \frac{u}{c^{(n)}}\right)}. \end{aligned}$$

The function $g_{n+1}(x, T)$ can be expressed as

$$\begin{aligned} g_{n+1}(x, T) = \mathbf{R}_n f_n(x, T) &= \frac{\bar{\mathbf{R}}_n f_n(x + m_n(T), T)}{Z_n(T)} + \frac{\frac{\partial}{\partial x} \bar{\mathbf{Q}}_n f_n(x + m_n(T), T)}{Z_n(T)} \frac{dm_n(T)}{dT} \\ &\quad - \frac{\bar{\mathbf{Q}}_n f_n(x + m_n(T), T)}{Z_n^2(T)} \frac{dZ_n(T)}{dT} \end{aligned}$$

with

$$Z_n(T) = \int_{-c^{(n)} M_n(T)}^{\infty} \bar{\mathbf{Q}}_n f_n(x, T) dx.$$

If the parameter $\kappa > 0$ in formula (2.13) $q_0(x, T)$ is sufficiently small and $n \leq N$, then the function $g_n(x, T)$ can be estimated similarly to the proof of Proposition 4.1 or Proposition 1 in [BM3]. Relation (A5) formulated below can be deduced from formula (A2) similarly to the proof of Lemma 1 of that paper. Then an argument similar to the proof of Lemma 2 in [BM3] enables one to prove formula (A4) formulated below. In this argument one can observe that a negligible error is committed if in the integrals appearing in the definition of $\bar{\mathbf{R}}_n f_n(x, T)$ the arguments $\ell_{n, M_n(T)}^\pm(x, u, v)$ defined in formula (2.25) are replaced by $\frac{x}{c_{n+1}} \pm u$. Some calculation also shows that we commit a

negligible error by replacing $\mathbf{R}_n f_n(x, T)$ with $\frac{\bar{\mathbf{R}}_n^{(1)} f_n(x, T)}{Z_n(T)}$. In such a way we get that

$$\begin{aligned} &\left| g_n(x, T) - \frac{\sqrt{A_0 - T}}{\sqrt{\pi}} \frac{2^{n/2}}{c^{(n)}} \left(x^2 - \frac{1}{2(A_0 - T)} \frac{c^{(n)^2}}{2^n} \right) \exp \left\{ -(A_0 - T) \frac{2^n x^2}{c^{(n)^2}} \right\} \right| \\ &\leq C(n) \kappa^{1/4} \exp \left\{ -\frac{(A_0 - T)}{4} \frac{2^n}{c^{(n)}} \left| 2x + \frac{x^2}{M_n^2(T)} \right| \right\} \quad \text{if } |x| < 2^{-n} \log \kappa^{-1}, \end{aligned} \tag{A4}$$

$$|g_n(x, T)| \leq C(n) \exp \left\{ -\frac{(A_0 - T)}{4} \frac{2^n}{c^{(n)}} \left| 2x + \frac{x^2}{M_n^2(T)} \right| \right\} \quad \text{for } x \geq -M_n(T), \quad (\text{A5})$$

$$|M_n(T) - M_0(T)| \leq C(n) \kappa^{1/2}, \quad \left| Z_n(t) - \frac{\sqrt{\pi}}{\sqrt{A_0 - t}} \right| \leq C(n) \kappa^{1/2}$$

with some constant $C(n)$ which may depend on n but not on the parameter κ of the model.

The previous results are sufficient to handle the functions $g_n(x, T)$ for small indices $n \leq N$. To work with indices $n \geq N$ we have to introduce, similarly to the argument in Section 4, the regularization of the functions $g_n(x, T)$, the linearization $\bar{\mathbf{U}}_n$ and $\mathbf{U}_n$ of the operators $\bar{\mathbf{R}}_n$ and $\mathbf{R}_n$ and to describe their action in the Fourier space.

Define the regularization of the function $g_n(x, T)$ as $\varphi_n(g_n(x, T)) = \frac{\partial \varphi_n(f_n(x, T))}{\partial T}$. We want to approximate the operator $\mathbf{R}_n$ with a simpler operator $\mathbf{U}_n$ in analogy with the approximation of $\mathbf{Q}_n$ by $\mathbf{T}_n$. Then we formulate and prove some inductive hypothesis about the behaviour of the operators $\mathbf{R}_n$ and $\mathbf{U}_n$.

A natural approximation of the operators $\bar{\mathbf{R}}_n$ and $\mathbf{R}_n$ by some operators $\bar{\mathbf{U}}_n$ and $\mathbf{U}_n$ can be obtained by differentiating $\bar{\mathbf{T}}_n \varphi(f_n(x, T))$ and $\mathbf{T}_n \varphi_n(f_n(x, T))$ with respect to the variable T . These considerations suggest the definition of the operators

$$\begin{aligned} \bar{\mathbf{U}}_n \varphi_n(f_n(x, T)) = & 2 \int e^{-v^2} \varphi_n \left(f_n \left(\frac{x}{c_{n+1}} + u + \frac{v^2}{2M_n(T)}, T \right) \right) \\ & \left\{ \varphi_n \left(g_n \left(\frac{x}{c_{n+1}} - u + \frac{v^2}{2M_n(T)}, T \right) \right) \right. \\ & \left. - v^2 \frac{M'_n(T)}{2M_n(T)^2} \frac{\partial}{\partial x} \varphi_n \left(f_n \left(\frac{x}{c_{n+1}} - u + \frac{v^2}{2M_n(T)}, T \right) \right) \right\} du dv \end{aligned}$$

with the function $g_n(x, T)$ defined in (A3) and

$$\mathbf{U}_n \varphi_n(f_n(x, T)) = \mathbf{U}_n^{(1)} \varphi_n(f_n(x, T)) + \mathbf{U}_n^{(2)} \varphi_n(f_n(x, T))$$

with

$$\begin{aligned} \mathbf{U}_n^{(1)} \varphi_n(f_n(x, T)) = & \frac{4}{c_{n+1} \sqrt{\pi}} \int e^{-v^2} \varphi_n \left(f_n \left(\frac{x}{c_{n+1}} + u - \frac{1}{4M_n(T)} + \frac{v^2}{2M_n(T)}, T \right) \right) \\ & \varphi_n \left(g_n \left(\frac{x}{c_{n+1}} - u - \frac{1}{4M_n(T)} - \frac{v^2}{2M_n(T)}, T \right) \right) du dv, \end{aligned}$$

and

$$\begin{aligned} \mathbf{U}_n^{(2)} \varphi_n(f_n(x, T)) = & \frac{4}{c_{n+1} \sqrt{\pi}} \int e^{-v^2} \left(\frac{M'_n(T)}{4M_n^2(T)} - v^2 \frac{M'_n(T)}{2M_n(T)^2} \right) \\ & \varphi_n \left(f_n \left(\frac{x}{c_{n+1}} - u - \frac{1}{4M_n(T)} + \frac{v^2}{2M_n(T)}, T \right) \right) \\ & \frac{\partial}{\partial x} \varphi_n \left(f_n \left(\frac{x}{c_{n+1}} - u - \frac{1}{4M_n(T)} + \frac{v^2}{2M_n(T)}, T \right) \right) du dv. \end{aligned}$$

The Fourier transform of $\tilde{\mathbf{U}}_n \varphi_n(f_n(x, T))$, $\mathbf{U}_n^{(1)} \varphi_n(f_n(x, T))$ and $\mathbf{U}_n^{(2)} \varphi_n(f_n(x, T))$ can be expressed as

$$\begin{aligned} \tilde{\mathbf{U}}_n \tilde{\varphi}_n(f_n(\xi, T)) &= c_{n+1} \sqrt{\pi} \frac{\tilde{\varphi}_n\left(f_n\left(\frac{c_{n+1}}{2}\xi, T\right)\right)}{\sqrt{1 + i \frac{c_{n+1}}{2M_n(T)}\xi}} \tilde{\varphi}_n\left(g_n\left(\frac{c_{n+1}}{2}\xi, T\right)\right) \\ &\quad - \frac{c_{n+1}^2 \sqrt{\pi}}{2} \frac{M'_n(T)}{M_n(T)^2} \xi \frac{\tilde{\varphi}_n^2\left(f_n\left(\frac{c_{n+1}}{2}\xi, T\right)\right)}{\left(1 + i \frac{c_{n+1}}{2M_n(T)}\xi\right)^{3/2}}, \\ \tilde{\mathbf{U}}_n^{(1)} \tilde{\varphi}_n(f_n(\xi, T)) &= 2 \frac{\exp\left\{\frac{ic_{n+1}\xi}{4M_n(T)}\right\}}{\sqrt{1 + i \frac{c_{n+1}}{2M_n(T)}\xi}} \tilde{\varphi}_n\left(f_n\left(\frac{c_{n+1}}{2}\xi, T\right)\right) \tilde{\varphi}_n\left(g_n\left(\frac{c_{n+1}}{2}\xi, T\right)\right) \quad (\text{A6}) \end{aligned}$$

and

$$\begin{aligned} \tilde{\mathbf{U}}_n^{(2)} \tilde{\varphi}_n(f_n(\xi, T)) &= \frac{c_{n+1} M'_n(T)}{4M_n(T)^2} \frac{\exp\left\{\frac{ic_{n+1}\xi}{4M_n(T)}\right\} \xi}{\sqrt{1 + i \frac{c_{n+1}}{2M_n(T)}\xi}} \\ &\quad \tilde{\varphi}_n^2\left(f_n\left(\frac{c_{n+1}}{2}\xi, T\right)\right) \left(1 - \frac{1}{1 + i \frac{c_{n+1}}{2M_n(T)}\xi}\right). \quad (\text{A7}) \end{aligned}$$

The above relation can also be extended to a larger set of the variables ξ in the complex plane by means of analytic continuation.

Now we formulate the inductive hypotheses we want to prove in the Appendix.

Property $K_1(n)$.

$$-\frac{dM_n(T)}{dT} > 0.$$

Property $K_2(n)$.

$$\begin{aligned} |g_n(x, T)| = \left| \frac{\partial}{\partial T} f_n(x, T) \right| &< K \left| \frac{dM_n(T)}{dT} \right| \exp \left\{ -\frac{1}{\sqrt{\beta_n(T)}} \left| 2x + \frac{x^2}{c^{(n)} M_n(T)} \right| \right\} \\ &\text{if } x > -c^{(n)} M_n(T) \end{aligned}$$

with a universal constant K .

Property $K_3(n)$.

$$|g_n(x, T) - \mathbf{U}_{n-1}\varphi_{n-1}(f_{n-1}(x, T))| < K \left| \frac{dM_n(T)}{dT} \right| \frac{\beta_n(T)}{c^{(n)}} \exp \left\{ -\frac{1.4}{\sqrt{\beta_n(T)}} \left| 2x + \frac{x^2}{c^{(n)}M_n(T)} \right| \right\} \quad \text{if } x > -c^{(n)}M_n(T)$$

with a universal constant K . The inequality remains valid if the function $g_n(x, T)$ is replaced by its regularization $\varphi_n(g_n(x, T))$.

The following property $K_4(n)$ which gives a bound on the Fourier transform of $\varphi_n(g_n(x, T))$ is an analog of Property $J(n)$.

Property $K_4(n)$.

$$|\tilde{\varphi}_n(g_n(-is, T))| = \left| \int e^{sx} \varphi_n(g_n(x, T)) dx \right| \leq \beta_n^{3/2}(T) s^2 \left| \frac{dM_n(T)}{dT} \right| e^{\beta_n(T)s^2} \quad \text{if } |s| < \frac{2}{\sqrt{\beta_{n+1}(T)}}.$$

In Property $K_4(n)$ we formulated a weaker estimate than in $J(n)$. It is enough to have a good bound on the moment generating function, i.e. on the analytic continuation of the Fourier transform to the imaginary axis together with the trivial estimate $|\tilde{\varphi}_n(g_n(-is + t, T))| \leq \tilde{\varphi}_n(g_n(-is, T))$ for all t .

The main result of the Appendix is the following Proposition A.

Proposition A. *Let the properties $K_1(m)$, $K_2(m)$, $K_3(m)$ and $K_4(m)$ hold in a neighbourhood of a parameter T together with the property $\frac{\beta_m(T)}{c^m} \leq \eta$ (with the same small number $\eta > 0$ which appeared in the proof of Propositions 4.1 and 4.2) for all $N \leq m \leq n$, and let also the inductive hypotheses $I(n)$ and $J(n)$ be also satisfied. Then the properties $K_1(n+1)$, $K_2(n+1)$, $K_3(n+1)$ and $K_4(n+1)$ also hold for this parameter T . The expression*

$$\delta_n(T) = \frac{d}{dT} \left(m_n(T) - \frac{1}{4M_n(T)} \right) = \frac{dm_n(T)}{dT} + \frac{1}{4M_n^2(T)} \frac{dM_n(T)}{dT},$$

satisfies the inequality

$$|\delta_n(T)| \leq C \left| \frac{dM_n(T)}{dT} \right| \frac{\beta_{n+1}(T)}{c^{(n+1)}} \beta_{n+1}(T) \quad (\text{A8})$$

with an appropriate $C > 0$, where $m_n(T)$ was defined in (2.22).

If we want to apply Proposition A, then first we have to show that properties $K_1(n)$, $K_2(n)$, $K_3(n)$ and $K_4(n)$ hold for $n = N$ if $T < c_0 A_0/2$. This can be done with

the help of an argument similar to the proof in the Corollary of Lemma 1 in [BM3]. Property $K_1(N)$ holds since $\frac{dM_N(T)}{dT}$ hardly differs from $\frac{dM_0(T)}{dT}$. Property $K_2(N)$ can be proved by means of relations (A4) and (A5). In the proof of Property $K_3(N)$ still the following additional observation is needed. Relation (A4) remains valid if the function $g_N(x, T) = \mathbf{R}_N f_{N-1}(x, T)$ is replaced by $\mathbf{U}_N \varphi_{N-1}(f_{N-1}(x, T))$ in this formula. (The term $\frac{dM_n(T)}{dT}$ on the right-hand side of the inductive hypotheses do not play an important role for $n = N$. It is strongly separated from zero if $T \leq c_0 A_0/2$.)

Relation $K_4(N)$ can be proved again with the help of formulas (A4), (A5) and the relations $\int \varphi_n(g_n(x, T)) dx = \int x \varphi_n(g_n(x, T)) dx = 0$. These relations imply that the value of the function $\tilde{\varphi}_n(g_n(s, T))$ and of its first derivative is zero in the point $s = 0$. Hence it is enough to give a good estimate of the second derivative of $\tilde{\varphi}_n(g_n(s, T))$.

Let us formulate the following Corollary of Proposition A.

Corollary. *Under the Conditions of Theorem 4 the set of the points T for which (n, T) is in the low temperature region is an interval $(0, T_n)$ for all $n \geq 0$. The inductive hypotheses $K_1(n)$, $K_2(n)$, $K_3(n)$ and $K_4(n)$ hold for all $T \in (0, T_n)$.*

Proof of the Corollary. The Corollary simply follows from Proposition A by means of induction with respect to n . In this induction we assume the statement of the Corollary for a fixed n together with the assumption that $\beta_n(T)$ is monotone increasing in the variable T for $0 < T < T_n$. The Corollary and the additional assumption hold for $n = N$ with $T_N = c_0 A_0/2$. If properties $K_1(n)$, $K_2(n)$, $K_3(n)$ and $K_4(n)$ hold for n , then because of Property $K_1(n)$ the function $M_n(T)$ is monotone decreasing and $\beta_{n+1}(T)$ is monotone increasing in the variable T . Then $T_{n+1} = \min(T_n, \max(T: \beta_{n+1}(T) < \eta))$, and by Proposition A the statements of the Corollary hold for $n + 1$.

Before turning to the proof of Proposition A we prove Theorem A with its help.

Proof of Theorem A. The proof of Part a.) is contained in the previous estimates of the Appendix. Part b.) can be obtained by differentiating the second formula in (2.24), and applying formula (A8).

Proof of Proposition A. Some calculation yields that because of properties $K_4(n)$, $J(n)$ relations (A6) and (A7) the Fourier transforms $\tilde{\mathbf{U}}_n^{(1)} \tilde{\varphi}_n(f_n(\xi, T))$, $\tilde{\mathbf{U}}_n^{(2)} \tilde{\varphi}_n(f_n(\xi, T))$ satisfy the inequalities

$$\begin{aligned} & \left| \tilde{\mathbf{U}}_n^{(1)} \tilde{\varphi}_n(f_n(t + is, T)) \right| \\ & \leq 2 \left| \frac{dM_n(T)}{dT} \right| \left(\frac{c_{n+1}}{2} s \right)^2 \beta_n(T)^{3/2} \exp \left\{ \left(\frac{c_{n+1}^2 \beta_n(T)}{2} + \frac{1}{M_n^2(T)} \right) s^2 \right\} \frac{1}{1 + \alpha_n(T) t^2} \end{aligned}$$

and

$$\begin{aligned} \left| \tilde{\mathbf{U}}_n^{(2)} \tilde{\varphi}_n(f_n(t + is, T)) \right| & \leq \frac{c_{n+1}^2 |M'_n(T)|}{8M_n(T)^3} (s^2 + t^2) \exp \left\{ \left(\frac{c_{n+1}^2 \beta_n(T)}{2} + \frac{1}{M_n^2(T)} \right) s^2 \right\} \\ & \frac{1}{(1 + \alpha_n(T) t^2)^2 \left(1 - \frac{c_{n+1}}{2M_n(T)} s \right)} \end{aligned}$$

for $|s| < \frac{4}{c_{n+1}\sqrt{\beta_{n+1}(T)}}$.

The function $\varphi_n(g_n(x), T)$ can be computed by means of the application of the inverse Fourier transformation and by replacement of the domain of integration from the real line to the line $\left\{ z = i \operatorname{sign} x \frac{2}{\sqrt{\beta_{n+1}(T)}} + t, t \in \mathbb{R}^1 \right\}$. We get, by applying the above estimates for the Fourier transforms $\tilde{\mathbf{U}}_n^{(1)}$ and $\tilde{\mathbf{U}}_n^{(2)}$ and exploiting the relation $\frac{M'_n(T)}{2M_n(T)^3} \leq \frac{1}{200}\beta_{n+1}(T)^2 \frac{dM_n(T)^2}{dT}$ together with the fact that the constants $\alpha_n(T)$ and $\beta_n(T)$ introduced in the definition of Properties $I(n)$ and $J(n)$ have the same order of magnitude that

$$\begin{aligned} |\mathbf{U}_n \varphi_n(f_n(x), T)| &\leq -K_1 \frac{dM_n(T)}{dT} e^{-2|x|\beta_{n+1}(T)^{-1/2}} \\ &\leq -K_2 \frac{dM_n(T)}{dT} \exp \left\{ -\frac{1}{\sqrt{\beta_{n+1}(T)}} \left| 2x + \frac{x^2}{c^{(n+1)}M_{n+1}(T)} \right| \right\}. \end{aligned} \quad (\text{A9})$$

The estimates obtained for $\tilde{\mathbf{U}}_n^{(1)}$ and $\tilde{\mathbf{U}}_n^{(2)}$ yield, with the choice $t = 0$ and some calculation that

$$\begin{aligned} \left| \tilde{\mathbf{U}}_n \tilde{\varphi}_n(f_n(-is, T)) \right| &\leq -\frac{9}{10} \frac{dM_n(T)}{dT} \beta_{n+1}(T)^{3/2} s^2 e^{\beta_{n+1}(T)s^2} \\ &\quad \text{if } |s| < \frac{2}{\sqrt{\beta_{n+2}(T)}}. \end{aligned} \quad (\text{A10})$$

(In the proof of Property $K_4(n+1)$ it will be important that the right-hand side of (A10) is less than the expression at the right-hand side of the formula which defines Property $K_4(n+1)$.)

We need a good estimate on the difference of $\mathbf{R}_n f_n(x, T) - \mathbf{U}_n \varphi_n(f_n(x, T))$ and its Fourier transform. These expressions can be bounded similarly to the proof of the corresponding inequalities in the proof of Proposition 3 in paper [BM3]. One has to compare the difference of the corresponding terms in the expressions $\mathbf{Q}_n \varphi_n(f_n(x, T))$ and $\mathbf{R}_n \varphi_n(f_n(x, T))$. Some calculation yields that

$$\left| Z_n(T) - \frac{c_{n+1}\sqrt{\pi}}{2} \right| \leq \frac{\beta_n(T)}{c^{(n)}}, \quad \left| m_n(T) + \frac{1}{4M_n(T)} \right| \leq \frac{\beta_n(T)}{c^{(n)}} \sqrt{\beta_n(T)}, \quad (\text{A11})$$

$$\begin{aligned} \left| \frac{dZ_n(T)}{dT} \right| &\leq -K \frac{\beta_n(T)}{c^{(n)}} \beta_n^{1/2}(T) \frac{dM_n(T)}{dT}, \\ \left| \frac{d}{dT} \left(m_n(T) + \frac{1}{4M_n(T)} \right) \right| &\leq -K \frac{\beta_{n+1}(T)}{c^{(n+1)}} \beta_{n+1}(T) \frac{dM_n(T)}{dT}. \end{aligned} \quad (\text{A12})$$

Relation (A8) is a consequence of (A12). Property $K_1(n+1)$ can be deduced from the above inequalities, since

$$\begin{aligned} -\frac{dM_{n+1}(T)}{dT} &= -\frac{dM_n(T)}{dT} + \frac{1}{c^{(n+1)}} \frac{dm_n(T)}{dT} \\ &\geq -\frac{dM_n(T)}{dT} \left(1 - \frac{1}{c^{(n+1)}} \left(\frac{1}{4M_n^2(T)} + K \frac{\beta_n^2(T)}{c^{(n)}} \right) \right) \geq -\frac{1}{2} \frac{dM_n(T)}{dT}. \end{aligned}$$

Now we turn to the proof of Property $K_3(n+1)$. We get, by applying again inequalities (A11) and (A12) together with the estimates obtained for $f_n(x, T)$, similarly to the proof of the estimates in the lemmas needed for the proof of Lemma 3 in [BM3] that

$$\begin{aligned} &\left| \frac{\bar{\mathbf{Q}}_n f_n(x + m_n(T), T)}{Z_n^2(T)} \frac{dZ_n(T)}{dT} \right| \\ &\leq K \frac{\beta_n(T)}{c^{(n)}} \left| \frac{dM_n(T)}{dT} \right| \exp \left\{ \frac{-1.5}{\sqrt{\beta_n(T)}} \left| 2x + \frac{x^2}{c^{(n)} M_n(T)} \right| \right\} \quad \text{if } x \geq c^{(n+1)} M_{n+1}(T), \end{aligned}$$

$$\begin{aligned} &\left| \frac{\frac{\partial}{\partial x} \bar{\mathbf{Q}}_n f_n(x + m_n(T), T)}{Z_n(T)} \frac{dm_n(T)}{dT} \right. \\ &\quad \left. - \frac{4}{c_{n+1} \sqrt{\pi}} \frac{M'_n(T)}{4M_n^2(T)} \int e^{-v^2} \varphi_n \left(f_n \left(\frac{x}{c_{n+1}} - u - \frac{1}{4M_n(T)} + \frac{v^2}{2M_n(T)}, T \right) \right) \right. \\ &\quad \left. \frac{\partial}{\partial x} \varphi_n \left(f_n \left(\frac{x}{c_{n+1}} - u - \frac{1}{4M_n(T)} + \frac{v^2}{2M_n(T)}, T \right) \right) du dv \right| \\ &\leq K \frac{\beta_n(T)}{c^{(n)}} \left| \frac{dM_n(T)}{dT} \right| \exp \left\{ \frac{-1.5}{\sqrt{\beta_n(T)}} \left| 2x + \frac{x^2}{c^{(n)} M_n(T)} \right| \right\} \quad \text{if } x \geq c^{(n+1)} M_{n+1}(T) \end{aligned}$$

and

$$\begin{aligned} &\left| \frac{\bar{\mathbf{R}}_n^{(2)} f_n(x + m_n(T), T)}{Z_n(T)} + \int v^2 e^{-v^2} \frac{M'_n(T)}{2M_n^2(T)} \right. \\ &\quad \left. \varphi_n \left(f_n \left(\frac{x}{c_{n+1}} + u - \frac{1}{4M_n(T)} + \frac{v^2}{2M_n(T)}, T \right) \right) \right. \\ &\quad \left. \frac{\partial}{\partial x} \varphi_n \left(f_n \left(\frac{x}{c_{n+1}} - u - \frac{1}{4M_n(T)} + \frac{v^2}{2M_n(T)}, T \right) \right) du dv \right| \\ &\leq -K \frac{\beta_n(T)}{c^{(n)}} \frac{dM_n(T)}{dT} \exp \left\{ -\frac{1.5}{\sqrt{\beta_n(T)}} \left| 2x + \frac{x^2}{c^{(n)} M_n(T)} \right| \right\} \end{aligned}$$

$$\text{if } x \geq c^{(n+1)} M_{n+1}(T).$$

To prove of Property $K_3(n+1)$ we need an estimate which compares the terms

$$\frac{\bar{\mathbf{R}}_n^{(1)} f_n(x + m_n(T), T)}{Z_n(T)} \quad \text{and} \quad \mathbf{U}_n^{(1)} \varphi_n(f_n(x, T)).$$

We claim that

$$\left| \frac{\bar{\mathbf{R}}_n^{(1)} f_n(x + m_n(T), T)}{Z_n(T)} - \mathbf{U}_n^{(1)} \varphi_n(f_n(x, T)) \right| \leq -K \frac{\beta_n(T)}{c^{(n)}} \frac{dM_n(T)}{dT} \\ \exp \left\{ -\frac{1.5}{\sqrt{\beta_n(T)}} \left| 2x + \frac{x^2}{c^{(n)} M_n(T)} \right| \right\} \quad \text{if } x \geq c^{(n+1)} M_{n+1}(T).$$

This estimate can be proved by means of Property $K_3(n)$. With the help of this relation it can be shown that a negligible error is committed if in the integrals defining $\bar{\mathbf{R}}_n^{(1)} f_n(x + m_n(T), T)$ and $\mathbf{U}_n^{(1)} \varphi_n(f_n(x, T))$ the functions g_n and $\varphi_n(g_n)$ are replaced by the function $\mathbf{U}_n \varphi_{n-1}(f_{n-1})$. After this replacement the proof of Theorem 3.2 can be adapted, since we can bound not only the function $\mathbf{U}_n \varphi_{n-1}(f_{n-1})$, but also its partial derivative with respect to the variable x .

These estimates together imply Property $K_3(n+1)$, and some calculation shows that a version of Property $K_3(n+1)$, where the function $g_{n+1}(x)$ is replaced by its regularization $\varphi_{n+1} g_{n+1}(x)$ is also valid. Since we gave a good estimate on $\mathbf{U}_n \varphi_n(f_n(x))$ in (A9), some calculation yields the proof of Property $K_2(n+1)$. It remained to prove Property $K_4(n+1)$.

Because of (A10) and (A12) (The latter formula together with (2.22 and (2.24) imply that formula (A10) remain valid with a slightly bigger coefficient if the term $\frac{dM_n(T)}{dT}$ is replaced by $\frac{dM_{n+1}(T)}{dT}$ in it), it is enough to give a good bound on the difference $\tilde{\varphi}_{n+1}(g_{n+1}(-is)) - \tilde{\mathbf{U}}_n \tilde{\varphi}_n(f_n(-is))$ to prove property $K_4(n+1)$. This can be done in the following way:

By applying the modified property of $K_3(n+1)$, where the function $g_{n+1}(x)$ is replaced by $\varphi_{n+1} g_{n+1}(x)$ we get that

$$\left| \frac{\partial^2}{\partial s^2} \left[\tilde{\varphi}_{n+1}(g_{n+1}(-is, T)) - \tilde{\mathbf{U}}_n \tilde{\varphi}_n(f_n(-is, T)) \right] \right| \\ \leq - \int x^2 \frac{dM_n}{dT} \frac{\beta_{n+1}(T)}{c^{(n+1)}} \exp \left\{ \left(|t| - \frac{2.8}{\sqrt{\beta_{n+1}(T)}} \right) x \right\} dx \leq K \frac{\beta_{n+1}^{5/2}(T)}{c^{(n+1)}} \frac{dM_n^2}{dT}$$

$$\text{if } |s| \leq \frac{2}{\sqrt{\beta_{n+2}(T)}}.$$

Since

$$\begin{aligned} & \tilde{\varphi}_{n+1}(g_{n+1}(0, T)) - \tilde{\mathbf{U}}_n \tilde{\varphi}_n(f_n(0, T)) \\ &= \frac{\partial}{\partial s} \left(\tilde{\varphi}_{n+1}(\tilde{g}_{n+1}(-is, T)) - \tilde{\mathbf{U}}_n \tilde{\varphi}_n(f_n(-is, T)) \right) \Big|_{s=0} = 0, \end{aligned}$$

the last relation implies that

$$\left| \tilde{\varphi}_{n+1}(\tilde{g}_{n+1}(-is, T)) - \tilde{\mathbf{U}}_n \tilde{\varphi}_n(f_n(-is, T)) \right| \leq -K \frac{\beta_{n+1}(T)}{c^{(n+1)}} \beta_{n+1}^{3/2} \frac{dM_n(T)}{dT} s^2$$

if $|s| \leq \frac{2}{\sqrt{\beta_{n+2}(T)}}$. This estimate together with relation (A10) imply Property $K_4(n+1)$ if the number η which is an upper bound for $\beta_{n+1}(T)/c^{(n+1)}$ is chosen sufficiently small. Theorem A is proved.

Appendix B. Proof of Proposition 1.2

Condition 1. We have that for $n \geq 1$,

$$1 < c_n = \left(\frac{1 + an}{1 + a(n-1)} \right)^\lambda$$

Observe that c_n is decreasing and

$$\lim_{n \rightarrow \infty} c_n = 1, \quad c_n \leq c_1 = (1 + a)^\lambda$$

This implies Condition 1.

Condition 2. We have that

$$(1 + an)^\lambda \sum_{j=n}^{n+K} (1 + aj)^{-\lambda} \geq \frac{K(1 + an)^\lambda}{(1 + a(n + K))^\lambda} \rightarrow K$$

as $n \rightarrow \infty$. This implies Condition 2.

Condition 3. For $k \leq n/2$ we estimate

$$l_k \sum_{j=k}^n l_j^{-1} = (1 + ak)^\lambda \sum_{j=k}^n (1 + aj)^{-\lambda} \geq C(1 + ak)^\lambda (1 + ak)^{-\lambda+1} = C(1 + ak)^{-1}$$

and for $k > n/2$ and $n \geq j \geq k$ we estimate

$$l_k l_j^{-1} \geq C_0 > 0$$

hence

$$l_k \sum_{j=k}^n l_j^{-1} \geq C_0(n-k+1)$$

Thus,

$$\sum_{k=1}^n \left(l_k \sum_{j=k}^n l_j^{-1} \right)^{-2} \leq C^{-2} \sum_{k=1}^{n/2} (1+ak)^{-2} + C_0^{-2} \sum_{k=n/2}^n (n-k+1)^{-2} \leq C_1$$

Condition 3 is checked.

Conditions 4 and 5 are obvious.

Acknowledgements. An essential part of this work was done at the Mathematisches Forschungsinstitut Oberwolfach, where the authors enjoyed their participation in the program “Research in Pairs”. They are thankful to the Mathematisches Forschungsinstitut for kind hospitality and the Volkswagen–Stiftung for support of their stay at Oberwolfach. The research of the first author (P.B.) was supported in part by the National Science Foundation, Grants No. DMS–9623214 and DMS–9970625, and this support is gratefully acknowledged.

References:

- [ACCN] M. Aizenman, J. T. Chayes, L. Chayes, C. M. Newman, Discontinuity of the magnetization in one-dimensional $1/|x-y|^2$ Ising and Potts models, *J. Statist. Phys.* **50**, 1–40 (1988).
- [AYH] P. W. Anderson, G. Yuval, and D. R. Hamann: *Phys. Rev.* **1B**, 1522 (1970).
- [BM1] P. M. Bleher and P. Major: Renormalization of Dyson’s hierarchical vector valued φ^4 model at low temperatures. *Comm. Math. Physics* **95**, 487–532 (1984).
- [BM2] P. M. Bleher and P. Major: Critical phenomena and universal exponents in statistical physics. On Dyson’s hierarchical model. *Annals of Probability* **15**, 431–477 (1987).
- [BM3] P. M. Bleher and P. Major: The large-scale limit of Dyson’s hierarchical vector-valued model at low temperatures. The non-Gaussian case. Part I. *Annales de l’Institut Henri Poincaré, Série Physique Théorique*, Volume **49** fascicule 1, 1–85 (1988).
- [BM4] P. M. Bleher and P. Major: The large-scale limit of Dyson’s hierarchical vector-valued model at low temperatures. The marginal case $c = \sqrt{2}$. *Comm. Math. Physics* **125**, 43–69 (1989).
- [Dys1] F. J. Dyson: Existence of a phase transition in a one-dimensional Ising ferromagnet, *Commun. Math. Phys.* **12**, 91–107 (1969).

- [Dys2] F. J. Dyson: An Ising ferromagnet with discontinuous long-range order, *Commun. Math. Phys.* **21**, 269–283 (1971).
- [Dys3] F. J. Dyson: Existence and nature of phase transitions in one-dimensional Ising ferromagnets. *Mathematical aspects of statistical mechanics (Proc. Sympos. Appl. Math., New York, 1971)*, pp. 1–12, SIAM-AMS Proceedings, Vol. V, *Amer. Math. Soc., Providence, R.I.*, (1972).
- [Fi] M. E. Fisher: *Rep. Progr. Phys.* **30**, 615–730 (1967).
- [Sim] B. Simon: Absence of continuous symmetry breaking in a one-dimensional n^{-2} model, *J. Statist. Phys.* **26**, 307–311 (1981).
- [Tho] D. J. Thouless: *Phys. Rev.* **187**, 732 (1969).
- [WK] K. G. Wilson and J. Kogut: The renormalization group and the ε -expansion, *Phys. Rep.* **12C**, 75–199 (1974).